

The Modified Red Flag Threat Index

Murdoch and Gitro (2010) introduced the Red Flag Threat Index (RFTI) as a simple index derived from expected relative humidity and wind speed values to quantify the severity of critical fire weather condition. The index was modeled after the Haines Index (Haines 1988), in that it used meteorological variables to rate fire danger independently of fuels. The Haines Index, however, assesses risk based on temperature-dewpoint depressions and lapse rates, and is thus primarily an indicator for plume-driven wildfires. In the Great Plains, relative humidity and wind speed are the most commonly used predictors for wind-driven grassland fires.

In 2011, Murdoch, Gitro, Lindley, and Vitale began work to make the RFTI more robust and to demonstrate the adaptability of the index for use in any climate regime subject to wind-driven fires in fine fuels. This effort involved statistical analyses of approximately 1,500 critical fire weather observations across west Texas and application of the index to a dataset of proximity observations for 99 significant wildfire starts within the West Texas Mesonet (WTM) domain (Lindley et al. 2011). The resultant Modified Red Flag Threat Index (RFTI Mod) provides a simple numerical and categorical rating which quantifies the threat posed by combinations of relative humidity and wind speed with respect to local climatological significance and past large wind-driven grassland wildfires. During the historic Texas wildfire season of 2011, the RFTI Mod was used with success by the Midland Fire Department as a strategic staffing-decision tool. They found that the use of the RFTI Mod decreased their overstaffing during false alarm days of marginal Red Flag-weather, and they credit use of the index in their decision to increase staff levels by 20 firefighters on 24 May 2011 – prior to the damaging CEED Fire which impacted the city of Midland.



The CEED Fire crossing Loop 250 into the western parts of Midland on 24 May 2011


The RFTI Mod and RFTI Mod Cat are now available in GFE and should be used instead of the previous RFTI and RFTI Cat. These values (which require the use of the RFTIMod_tool and RFTIModCat_tool) will be added to the PFWLUB for use by local decision makers.

Index Derivation and Application –


Just as with the original RFTI, the RFTI Mod is derived as the simple summation of a *relative humidity term RFTI(A)* and a *wind speed term RFTI(B)*.

$$\text{RFTI Mod} = \text{RFTI(A)} + \text{RFTI(B)}$$


The distinguishing factor between the original RFTI and the RFTI Mod is that for the RFTI Mod these terms are based on statistical quartile rankings for a 10-year climatological database of critical fire weather observations, or observations exceeding the local Red Flag Warning criteria, at the location of interest. During the development of the RFTI Mod, such climatological thresholds of 2-m relative humidity and 6-m wind speed were found at Amarillo (KAMA), Lubbock (KLBB), and Midland (KMAF) spanning 2000-2009. Note, 10-m observed wind speeds were reduced to 6-m via a 0.886 factor as observed by the WTM at Reese Center during diurnal burn periods (Lindley et al. 2011).



climatological quartile rankings and RFTI(A) and RFTI(B) terms at Amarillo (KAMA)




climatological quartile rankings and RFTI(A) and RFTI(B) terms at Lubbock (KLBB)




climatological quartile rankings and RFTI(A) and RFTI(B) terms at Midland (KMAF)

In order to investigate the RFTI Mod's relevance to west Texas wildfire environments, the index was applied to the Lindley et al. (2011) dataset of 99 proximity meteorological observations for significant wind-driven grassland wildfire starts within the WTM domain. The dataset composed of wildland fire starts that grew to consume burn areas ≥ 300 acres spanning the 2006-2010 fire seasons, was differentiated spatially so that each fire start would be compared to the appropriate climatological thresholds for the nearest site, either KAMA, KLUB, or KMAF.




Lindley et al. (2011) wildfire start dataset shown as applied to the appropriate climatological quartiles for critical fire weather at KAMA, KLBB, and KMAF

Application of the Lindley et al. 2011 significant fire start dataset to the climatological quartiles for critical fire weather across west Texas reveals several useful applications. First, the number of significant fire starts per RFTI Mod rating approximates a normal distribution centered on the Critical-Low and Critical-High RFTI Mod values of 3-6. Secondly, the average fire size in each RFTI Mod category generally increases with increasing RFTI Mod values, and all fires that grew to consume > 35,000 acres occurred with RFTI Mod values ≥ 5 . Also, only one fire of 640 acres occurred with a RFTI Mod of 0, and no fires occurred in environments that exceeded the 10-year climate database of either 2-m relative humidity and 6-m wind speed, however, such high-end values should not be discounted and corresponding RFTI Mod values of 9 and 10 have been assigned the categorical rating “Historically Critical”.



Application of the Lindley et al. (2011) wildfire start dataset to 10-year climatological quartiles for critical fire weather across west Texas

For use in this document, the Lubbock (KLBB) RFTI Mod was applied to the February-May 2011 Southern Plains wildfire outbreaks. All of the outbreak days had a maximum RFTI Mod ≥ 5 (Critical-High). Arguably the days with the most significant impacts to the Lubbock-area proper, 27 February (central Lubbock fire) and 24 May (Reese Center fire) correspond to RFTI Mod of 8 (Extremely Critical).



Lubbock (KLBB) RFTI Mod values for the February-May 2011 Southern Plains wildfire outbreaks

Conclusion –

The modified version of the Red Flag Threat Index (RFTI Mod) shows operational value as a decision tool which quantifies the severity of critical fire weather relative to a statistical analysis of climatological critical fire weather independent of vegetative fuels. The RFTI Mod is derived by scoring the statistical quartiles of approximately 1,500 critical fire weather observations across west Texas for 2-m relative humidity and 6-m wind speed in excess of local Red Flag Warning criteria. When applied to a pre-existing dataset of WTM proximity meteorological observations for 99 significant wind-driven wildfire starts, the distribution of fire start environments approximates a normal distribution relative to RFTI Mod centered on values of 3-6 (Critical-Low and Critical-High). This distribution of RFTI Mod scores shows minimal fire activity when neither 2-m relative humidity nor 6-m wind speed exceed critical thresholds (RFTI Mod=0), and as expected, no fires within the database occurred with either 2-m relative humidity or 6-m wind speeds in excess of a 10-year climatology. Such rare and extreme values, however, should not be discounted and would be considered “Historically Critical”. All fires that consumed burn areas $\geq 35,000$ acres were associated with RFTI Mod values ≥ 5 . Also, each of the February – May 2011 Southern Plains wildfire outbreaks occurred during RFTI Mod values ≥ 5 at Lubbock (KLBB). The outbreak days with the most significant local impacts at Lubbock occurred when the KLBB RFTI Mod reached 8 (Extremely-Critical).

References –

Haines, DA, 1988: A Lower Atmosphere Severity Index for Wildland Fires. *National Weather Digest* 13, 23-27.

Lindley, T. T., J. D. Vitale, W. S. Burgett, and M.-J. Beierle, 2011: Proximity meteorological observations for wind-driven grassland wildfire starts on the southern High Plains. *Electronic J. Severe Storms Meteor.*, 6(1), 1-27

Murdoch, G. P., C. Gitro, 2010: Assessing critical fire weather conditions using a red flag threat index. 35th Annual Nat. Weath. Assoc. Meeting. Tucson, AZ.

Contact Information –

Greg Murdoch – NOAA/NWS Midland, Texas Gregory.Murdoch@noaa.gov

Todd Lindley – NOAA/NWS Lubbock, Texas Todd.Lindley@noaa.gov

