## Red River and Devils Lake Basin - 2022 Spring Flood Outlook



Discussion Points 2/10/2022 prepared by




NWS - Weather Forecast Office, Grand Forks ND NWS - North Central River Forecast Center, Chanhassen MN

This outlook is for the U.S. portion of the basin and is based on conditions through Tuesday, 2/08/2022. All graphics, probabilities, and related discussions are available at **weather.gov/fgf**. The next update will be issued by 2/24/2022.

## **Bottom Line up Front!**

- Its mid-winter... and there's a moderate to somewhat high threat for *significant* snowmelt flooding:
  - Lingering 2021 drought conditions mean that surface storage has room for snowmelt infiltration - if thaw is gentle.
  - Snowfall/SWE through early January is near to above long term mid-winter normal amounts.
  - Frost depths are running near to somewhat deeper than long term normal, due to cold January and early February conditions.
- A turnaround from record driest year in 2021. But, not near as wet as the recent wet years of 2019 to early 2020. Soil

moisture is back to within an inch plus-or-minus of normal soil moisture.



- Climate outlooks currently indicate a trend change in February, from cooler and snowier than normal towards near normal temperatures and precipitation through early spring, which helps to keep our overall risk in check.

Long Story Short: The risk for significant snowmelt flooding is moderately high, running near to slightly above long-term historical averages across the Red River and Devils Lake Basins (U.S. portions).

## **Key Snowmelt Flood Components:**

- **1. Base Streamflow:** Near normal for this time of year. USGS analyses indicate that the Red River and most of its ND and MN tributaries are moderate ice covered and/or flowing at 25% -75% of normal ranges. Somewhat higher in the far southern basin and somewhat lower near the CanAm border [link: https://waterdata.usgs.gov/nwis/rt].
- 2. Soil Moisture at Freeze-up: Near normal. From slightly above in the far south to slightly below in the far north basins. [Link: https://www.cpc.ncep.noaa.gov/products/Soilmst Monitoring/US/Soilmst/Soilmst.shtml]
- **3. Frost Depth:** Near to slightly Deeper than normal. A quite cold mid-winter period has allowed for deep frost penetration in most areas. Frost depths range from 12 to 40 inches in most locations, with deepest frost penetration across the northern reaches, and in upland areas of northwest MN, where early snowpack was lighter. Lake/River ice thicknesses are running near long term normal and is quite variable. [Link: https://www.weather.gov/ncrfc/LMI\_FrostDepthMap]
- **4. Winter Snowpack/SWE: Near to Above normal.** Since Dec 1<sup>st</sup>, snowfall runs from 90-130 percent of normal, SWE ranges from 2.0 to 4.0 inches. Fairly well distributed across the area, though quite splotchy due to relocation of snow during blowing snow and blizzard episodes. [Link: <a href="https://www.nohrsc.noaa.gov/nsa/">https://www.nohrsc.noaa.gov/nsa/</a>]
- **5.** Along with our flood partners, we've developed a display graphic which relates the current flood outlook to our historical flood levels, now available for all our forecast locations! *Check it out at:* https://www.weather.gov/fgf/PFOS

DEVILS LAKE & SI			Long bruary				
OCATION	95%	90%	75%	50%	25%	10%	05%
 CREEL BAY			1450 2				
CREEL BAY EAST STUMP LAKE	1449.8 1449.8						
he current heights	of Devi	ls Lake	and St	ump Lak	e are ~	1447.28	ft. M
Color code: Below	w Min	nor M	oderate	Maj	or F	lood of	Record
RED RIVER AND TR			Long lid Feb	_			
OCATION			75%				
WAHPETON	11.6		12.0				
HICKSON							
FARGO	24.8 <b>24.0</b>	25.2	27.2	30.9	33.9	35.9	38.3
	23.6	24.7	28.7	32.9	36.6	38.7	39.1
GRAND FORKS							
OSLO	34.0	34.4	35.3	36.3	37.1	38.4	39.0
DRAYTON	<b>34.0</b> 36.8	38.0	39.8	41.1	42.2	42.9	43.6
PEMBINA							
	13.4	10.5			Tribut		55.0
South Fork Buffalo							
	13.4	13.9	14.5	15.0	15.7	16.5	17.6
Buffalo River	6 0	<i>6</i> F	7 2	0 0	0.2	10 1	10 5
HAWLEY DILWORTH	6.0	0.5	7.3 <b>18.5</b>	8.0	9.3	10.1	T0.5
		16.4	18.5	20.1	21.6	22.8	23.8
Wild Rice River		<i>c</i>			0 0	40.0	44.0
TWIN VALLEY	5.7	6.0	6.6 <b>24.4</b>	7.6	9.2	10.2	11.2
HENDRUM	19.9	21.9	24.4	28.1	30.3	31.6	32.0
Marsh River		9.6	10.7	12.3	15.6	17.2	18.6
Sand Hill River CLIMAX		15 Ω	18.5	22 8	26 1	30 1	31 2
Red Lake River		13.0	10.0	22.0	20.1	30.1	51.2
HIGH LANDING		7.1	8.3	9.4	10.9	11.8	12.1
CROOKSTON			18.3				31.0
Snake River			<i>-</i> -	·			
ABOVE WARREN	63.7	63.9	64.2	65.1	65.9	67.0	68.5
ALVARADO Two Rivers River	101.3	101.9	103.0	105.3	107.4	108.6	109.3
HALLOCK	804.1	804.6	805.8	807.5	808.6	809.4	810.4
Roseau River ROSEAU	11.7	12.1	13.3	14.6	15.6	17.4	18.1
	North Dakota Tributaries:						
Wild Rice River ABERCROMBIE		11 0	14.1	16.0	19.8	23.2	24.6
Shevenne River							
VALLEY CITY LISBON	12.2	12.6	13.4	15.8	19.2	22.1	23.0
LISBON	12.6	13.1	14.6	16.0	19.4	22.5	24.1
KINDRED	15.9	16.8	18.5	20.3	21.2	21.2	21.2
WEST FARGO DVRSN	15.7	17.2	18.0	21.1	21.3	21.3	21.3
HARWOOD Maple River			87.9				
ENDERLIN	11 2	11 6	12.2	12 7	12 7	14.5	15.2
MAPLETON			20.5				
Goose River				,			
HILLSBORO	10.4	11.4	13.1	13.8	14.9	15.8	16.4
Forest River MINTO	5.1	5.4	5.8	6.8	7.2	7.8	7.9
Park River GRAFTON*							
Pembina River							
WALHALLA	5 3	5.5	6.6	8 0	10 1	13.0	13.8
MATITATIA							

## Notes

1. Devils Lake Basin Runoff Risk is moderately high. An additional rise of 3 to 4 feet is expected (75% to 25% risk range). A slight increase from the Jan 27<sup>th</sup> Outlook. A ½ to 1 ft. rise on Devils Lake is considered about normal.

Note: Devils Lake is currently about 1.25 feet lower than this time last year.

- 2. Red River Basin Flood Risk is moderately high. All Red River main-stem points may see near to somewhat above normal spring runoff/flows.
- near to above normal soil moistures in the south, near normal frost depths.
- coupled with moderate to high winter snowpack and SWE to date.
- 3. Near normal snowpack and runoff potential is evident in most all MN tributaries.

Near to slightly below normal soil moisture and near normal snowpack throughout westcentral and northwest MN.

4. ND tributaries have a slightly higher runoff potential across southeast ND that decreases as one moves north into northeast ND.

Mid and Upper Sheyenne basin soils are near normal moisture, while snowpack is somewhat above normal.

Lower Sheyenne through eastcentral ND tribs have near normal soil moisture and near to above normal snowpack.

Northeast ND has slightly drier soils with near normal snowpack. All values up slightly since Jan 27<sup>th</sup> Outlook.

Note: new Grafton Bypass!