

Model Output Statistics (MOS) - Objective Interpretation of NWP Model Output


University of Maryland – April 4, 2012

Mark S. Antolik

Meteorological Development Laboratory
Statistical Modeling Branch
NOAA/National Weather Service
Silver Spring, MD

(301) 713-0023 ext. 110

email: mark.antolik@noaa.gov



MOS Operational System “Fun Facts”

With apologies to David Letterman,
of course!



- 9 million regression equations
- 75 million forecasts per day
- 1200 products sent daily
- 400,000 lines of code – mostly FORTRAN
- 180 min. supercomputer time daily
- All developed and maintained by ~ ~~12~~⁸ MDL / SMB meteorologists!



OUTLINE

1. Why objective statistical guidance?

2. What is MOS?

Definition and characteristics

The “traditional” MOS product suite (GFS, NAM)

Other additions to the lineup

3. Simple regression examples / REEP

4. Development strategy -

MOS in the “real world”

5. Verification

6. Dealing with NWP model changes

7. Where we’re going – GMOS and the future

WHY STATISTICAL GUIDANCE?

- **Add value to direct NWP model output**

Objectively interpret model

- **remove systematic biases**
- **quantify uncertainty**

Predict what the model does not

Produce site-specific forecasts

(i.e. a “downscaling” technique)


- **Assist forecasters**

“First Guess” for expected local conditions


“Built-in” model/climo memory for new staff

A SIMPLE STATISTICAL MODEL

Relative Frequency of Precipitation as a Function of
12-24 Hour NGM Model-Forecast Mean RH



MOS Max Temp vs. Direct Model Output



What is MOS?

MODEL OUTPUT STATISTICS (MOS)

Relates observed weather elements (**PREDICTANDS**) to appropriate variables (**PREDICTORS**) via a statistical approach.

Predictors are obtained from:

1. Numerical Weather Prediction (NWP) Model Forecasts
2. Prior Surface Weather Observations
3. Geoclimatic Information

Current Statistical Method:

MULTIPLE LINEAR REGRESSION
(Forward Selection)

MODEL OUTPUT STATISTICS (MOS)

Properties

- **Mathematically simple, yet powerful**
- **Need historical record of observations at forecast points**
(Hopefully a long, stable one!)
- **Equations are applied to future run of similar forecast model**

MODEL OUTPUT STATISTICS (MOS)

Properties (cont.)

- **Non-linearity can be modeled by using NWP variables and transformations**
- **Probability forecasts possible from a single run of NWP model**
- **Other statistical methods can be used e.g. Polynomial or logistic regression; Neural networks**

MODEL OUTPUT STATISTICS (MOS)

- **ADVANTAGES**

- Recognition of model predictability

- Removal of some systematic model bias

- Optimal predictor selection

- Reliable probabilities

- Specific element and site forecasts

- **DISADVANTAGES**


- Short samples

- Changing NWP models

- Availability & quality of observations

MAJOR CHALLENGE TO MOS DEVELOPMENT:

RAPIDLY EVOLVING NWP MODELS AND OBSERVATION PLATFORMS



Can make for:

1. SHORT, UNREPRESENTATIVE DATA SAMPLES
2. DIFFICULT COLLECTION OF APPROPRIATE PREDICTAND DATA

New observing systems: (ASOS, WSR-88D, Satellite)
(Co-Op, Mesonets)

“Old” predictands: The elements don’t change!

“Traditional” MOS text products

GFS MOS GUIDANCE MESSAGE

FOUS21-26 (MAV)

| KLNS | GFS MOS GUIDANCE | | | | | | | | | | | | | | | | | | | 11/29/2004 | | | | 1200 UTC | | | |
|------|------------------|----|----|----|------|----|--------|----|----|--------|----|----|----|----|------|----|----|----|-------|------------|------|---|----|----------|--|--|--|
| DT | /NOV 29/NOV | | | 30 | | | /DEC 1 | | | /DEC 2 | | | | | | | | | | | | | | | | | |
| HR | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 06 | 12 | | | | | | |
| N/X | | | | | | | 28 | | | | 48 | | | | 35 | | | 49 | | 33 | | | | | | | |
| TMP | 43 | 44 | 39 | 36 | 33 | 32 | 31 | 39 | 46 | 45 | 41 | 38 | 37 | 39 | 41 | 44 | 45 | 44 | 40 | 40 | 35 | | | | | | |
| DPT | 27 | 27 | 28 | 29 | 29 | 29 | 29 | 33 | 35 | 35 | 36 | 35 | 36 | 39 | 41 | 42 | 37 | 34 | 30 | 30 | 28 | | | | | | |
| CLD | CL | BK | BK | BK | OV | OV | OV | OV | OV | OV | OV | OV | OV | OV | OV | OV | OV | BK | CL | CL | CL | | | | | | |
| WDR | 34 | 36 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 14 | 12 | 12 | 10 | 11 | 12 | 19 | 28 | 29 | 29 | 29 | 28 | | | | | | |
| WSP | 06 | 02 | 00 | 00 | 00 | 00 | 00 | 00 | 00 | 01 | 02 | 04 | 04 | 06 | 07 | 08 | 15 | 17 | 18 | 09 | 05 | | | | | | |
| P06 | | | 0 | | 0 | | 4 | | 3 | | 11 | | 65 | | 94 | | 96 | | 7 | 0 | 0 | | | | | | |
| P12 | | | | | | | 6 | | | | 19 | | | | 94 | | | | 96 | | 0 | | | | | | |
| Q06 | | | 0 | | 0 | | 0 | | 0 | | 0 | | 3 | | 4 | | 4 | | 0 | 0 | 0 | | | | | | |
| Q12 | | | | | | | 0 | | | | 0 | | | | 4 | | | | 2 | | 0 | | | | | | |
| T06 | | 0/ | 0 | | 0/18 | | 0/ | 3 | | 0/ | 0 | | 0/ | 0 | 0/18 | | 2/ | 1 | 10/ | 4 | 0/ | 3 | 1/ | 0 | | | |
| T12 | | | | | 0/26 | | | | | 0/17 | | | | | 0/27 | | | | 10/25 | | 1/38 | | | | | | |
| POZ | 2 | 0 | 0 | 1 | 2 | 4 | 4 | 0 | 1 | 1 | 2 | 3 | 3 | 1 | 1 | 0 | 2 | 1 | 2 | 3 | 1 | | | | | | |
| POS | 13 | 2 | 1 | 2 | 1 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 2 | 0 | 0 | 0 | 3 | 0 | 9 | 28 | | | | | | |
| TYP | R | R | R | R | R | R | R | R | R | R | R | R | R | R | R | R | R | R | R | R | R | | | | | | |
| SNW | | | | | | | 0 | | | | | | | | 0 | | | | | | 0 | | | | | | |
| CIG | 8 | 8 | 8 | 8 | 7 | 7 | 7 | 8 | 8 | 7 | 7 | 7 | 4 | 2 | 3 | 3 | 6 | 7 | 8 | 8 | 8 | | | | | | |
| VIS | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 5 | 5 | 4 | 2 | 6 | 7 | 7 | 7 | 7 | | | | | | |
| OBV | N | N | N | N | N | N | N | N | N | N | N | N | BR | BR | BR | BR | N | N | N | N | N | | | | | | |

NAM MOS GUIDANCE MESSAGE

FOUS44-49 (MET)

| KBWI | NAM MOS GUIDANCE | | | | | | | | | | | | | | | | | | | 2/27/2009 | | | | 1200 UTC | | | |
|------|------------------|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|----|--------|-----------|----|--------|--|----------|--|--|--|
| DT | /FEB 27/FEB | | | 28 | | | | | | | | | | | | | | | /MAR 1 | | | /MAR 2 | | | | | |
| HR | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 06 | 12 | | | | | | |
| N/X | | | | | | | 38 | | | | 46 | | | | 32 | | | 41 | | 24 | | | | | | | |
| TMP | 59 | 58 | 55 | 54 | 49 | 43 | 38 | 38 | 43 | 45 | 40 | 38 | 37 | 35 | 33 | 34 | 37 | 38 | 33 | 29 | 25 | | | | | | |
| DPT | 46 | 47 | 48 | 46 | 37 | 30 | 24 | 22 | 22 | 22 | 24 | 27 | 28 | 26 | 25 | 24 | 24 | 21 | 17 | 12 | 10 | | | | | | |
| CLD | OV | OV | OV | OV | OV | SC | SC | SC | CL | BK | OV | OV | OV | OV | OV | OV | OV | OV | OV | OV | BK | | | | | | |
| WDR | 21 | 20 | 22 | 25 | 31 | 32 | 34 | 36 | 01 | 03 | 05 | 04 | 01 | 36 | 35 | 35 | 35 | 34 | 35 | 33 | 34 | | | | | | |
| WSP | 15 | 09 | 08 | 06 | 10 | 11 | 10 | 12 | 10 | 09 | 08 | 10 | 12 | 13 | 14 | 16 | 11 | 13 | 15 | 16 | 17 | | | | | | |
| P06 | | | 89 | | 10 | | 3 | | 2 | | 2 | | 76 | | 73 | | 13 | | 17 | 27 | 19 | | | | | | |
| P12 | | | | | | | 10 | | | | 3 | | | | 81 | | | 17 | | 30 | | | | | | | |
| Q06 | | | 1 | | 0 | | 0 | | 0 | | 0 | | 4 | | 1 | | 0 | | 0 | 0 | 0 | | | | | | |
| Q12 | | | | | | | 0 | | | | 0 | | | | 4 | | | 0 | | 0 | | | | | | | |
| T06 | | 2/ | 9 | 0/ | 5 | 0 | /0 | 0/ | 5 | 3/ | 1 | 5/ | 3 | 0/ | 0 | 0/ | 2 | 2/ | 5 | 0/ | 0 | | | | | | |
| T12 | | | | 2/ | 9 | | | 0/ | 5 | | | 5/ | 3 | | | 1/ | 2 | | 7/ | 5 | | | | | | | |
| SNW | | | | | | | 0 | | | | | | | | 0 | | | | | | 0 | | | | | | |
| CIG | 6 | 6 | 4 | 5 | 7 | 8 | 8 | 8 | 8 | 8 | 7 | 6 | 4 | 3 | 4 | 3 | 4 | 4 | 7 | 6 | 7 | | | | | | |
| VIS | 7 | 7 | 6 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 3 | 6 | 5 | 7 | 7 | 7 | 7 | 7 | 7 | | | | | | |
| OBV | N | N | N | N | N | N | N | N | N | N | N | N | BR | N | BR | N | N | N | N | N | N | | | | | | |

Short-range (GFS / NAM) MOS

- **STATIONS:**

- Now at approx. 1990 Forecast Sites (CONUS, AK, HI, PR, Canada)

- **FORECASTS:**





- Available at projections of 6-84 hours
GFS available for 0600 and 1800 UTC cycles

- **RESOLUTION:**


- GFS predictors on 95.25 km grid; NAM on 32 km
Predictor fields available at 3-h timesteps

- **DEPENDENT SAMPLE NOT “IDEAL”:**

- Fewer seasons than older MOS systems
Non-static underlying NWP model



**Approx.
1990 sites**



MOS Stations

Short-range (GFS / NAM) MOS

- **STATIONS:**

- Now at approx. 1990 Forecast Sites (CONUS, AK, HI, PR)

- **FORECASTS:**

- Available at projections of 6-84 hours
GFS available for 0600 and 1800 UTC cycles

- **RESOLUTION:**

- GFS predictors on 95.25 km grid; NAM on 32 km
Predictor fields available at 3-h timesteps

- **DEPENDENT SAMPLE NOT “IDEAL”:**

- Fewer seasons than older MOS systems
Non-static underlying NWP model

GFSX MOS GUIDANCE MESSAGE

FEUS21-26 (MEX)

| KCXY | GFSX MOS GUIDANCE 11/26/2004 0000 UTC | | | | | | | | | | | | | | | | | | |
|------|---------------------------------------|-----|----|-----|-----|-----|----|-----|-----|-----|-----|-----|-----|-----|-----|-------|----|--|--|
| FHR | 24 | 36 | 48 | 60 | 72 | 84 | 96 | 108 | 120 | 132 | 144 | 156 | 168 | 180 | 192 | | | | |
| FRI | 26 | SAT | 27 | SUN | 28 | MON | 29 | TUE | 30 | WED | 01 | THU | 02 | FRI | 03 | CLIMO | | | |
| X/N | 43 | 29 | 47 | 40 | 55 | 35 | 51 | 29 | 45 | 32 | 40 | 36 | 42 | 30 | 45 | 31 | 46 | | |
| TMP | 37 | 32 | 43 | 43 | 46 | 37 | 41 | 32 | 39 | 35 | 36 | 38 | 37 | 33 | 37 | | | | |
| DPT | 24 | 27 | 37 | 40 | 32 | 28 | 28 | 26 | 31 | 32 | 30 | 32 | 27 | 24 | 25 | | | | |
| CLD | PC | OV | OV | OV | PC | CL | PC | PC | OV | OV | OV | PC | CL | CL | CL | | | | |
| WND | 10 | 5 | 11 | 11 | 16 | 10 | 10 | 5 | 9 | 6 | 10 | 12 | 14 | 12 | 12 | | | | |
| P12 | 0 | 5 | 13 | 91 | 13 | 3 | 9 | 14 | 24 | 52 | 54 | 48 | 21 | 12 | 25 | 20 | 18 | | |
| P24 | | | 16 | | 100 | | 9 | | 26 | | 62 | | 72 | | 25 | | 29 | | |
| Q12 | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 0 | 2 | 2 | 2 | | | | | | | |
| Q24 | | | 0 | | 3 | | 0 | | 0 | | 4 | | | | | | | | |
| T12 | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 4 | 6 | 4 | 3 | 1 | 1 | 1 | | | | |
| T24 | | 0 | | 3 | | 0 | | 0 | | 6 | | 4 | | 1 | | | | | |
| PZP | 12 | 9 | 12 | 4 | 3 | 5 | 6 | 10 | 8 | 8 | 3 | 16 | 10 | 12 | 8 | | | | |
| PSN | 62 | 15 | 3 | 0 | 0 | 10 | 9 | 15 | 24 | 1 | 0 | 9 | 32 | 27 | 18 | | | | |
| PRS | 26 | 24 | 7 | 0 | 17 | 18 | 20 | 13 | 15 | 1 | 2 | 18 | 9 | 11 | 11 | | | | |
| TYP | S | RS | R | R | R | R | R | R | RS | R | R | R | RS | RS | R | | | | |
| SNW | | | 0 | | 0 | | 0 | | 0 | | | | | | | | | | |



**MOS station-oriented products:
Other additions**

Marine MOS

44004 GFS MOS GUIDANCE 11/22/2005 1200 UTC

| DT | /NOV 22/NOV 23 | | | | | | /NOV 24 | | | | | | /NOV 25 | | | | | | | | |
|------|----------------|----|----|----|----|----|---------|----|----|----|----|----|---------|----|----|----|----|----|----|----|----|
| HR | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 03 | 06 |
| TMP | 58 | 53 | 49 | 49 | 50 | 48 | 46 | 44 | 44 | 45 | 47 | 48 | 51 | 54 | 56 | 60 | 62 | 61 | 59 | 51 | 47 |
| WD | 23 | 25 | 27 | 28 | 28 | 29 | 29 | 28 | 28 | 27 | 27 | 25 | 22 | 22 | 22 | 23 | 23 | 23 | 24 | 27 | 28 |
| WS | 33 | 31 | 29 | 25 | 23 | 22 | 24 | 25 | 23 | 18 | 14 | 12 | 14 | 19 | 26 | 29 | 30 | 29 | 29 | 28 | 24 |
| WS10 | 36 | 34 | 31 | 26 | 25 | 24 | 26 | 27 | 25 | 19 | 15 | 13 | 15 | 21 | 28 | 31 | 32 | 31 | 31 | 30 | 26 |

| DT | /NOV 25 | | | | | | / |
|------|---------|----|----|----|----|----|---|
| HR | 09 | 12 | 15 | 18 | 21 | 00 | |
| TMP | 45 | 45 | 45 | 47 | 47 | 47 | |
| WD | 29 | 29 | 28 | 30 | 29 | 34 | |
| WS | 18 | 15 | 10 | 10 | 13 | 12 | |
| WS10 | 20 | 16 | 11 | 11 | 14 | 13 | |





Max/Min Guidance for Co-op Sites

GFS-BASED MOS COOP MAX/MIN GUIDANCE 3/01/05 1800 UTC

| | WED 02 | THU 03 | FRI 04 | |
|--------------|----------------|----------------|----------------|-------------------------|
| ANNM2 | 26 46 | 24 45 | 25 46 | |
| BERM2 | 28 41 | 25 39 | 25 43 | |
| BTVM2 | 23 39 | 21 38 | 20 43 | ← Beltsville, MD |
| CBLM2 | 20 40 | 18 39 | 20 46 | |
| CHEM2 | 25 42 | 21 39 | 21 44 | |
| CNWM2 | 21 42 | 21 40 | 20 45 | |
| DMAM2 | 20 37 | 18 37 | 20 42 | |
| ELCM2 | 25 41 | 21 41 | 18 45 | |
| EMMM2 | 23 42 | 20 41 | 20 43 | |
| FREM2 | 23 46 | 21 42 | 23 44 | |
| FRSM2 | 17 27 | 13 27 | 13 36 | |
| GLDM2 | 21 37 | 18 39 | 18 43 | ← Glenn Dale, MD |
| HAGM2 | 23 43 | 18 43 | 19 45 | |
| KAPG | 27 41 | 23 37 | 22 43 | |
| LRLM2 | 23 44 | 21 42 | 22 46 | ← Laurel 3 W |
| MECM2 | 24 47 | 20 42 | 20 45 | |
| MILM2 | 25 48 | 22 41 | 20 39 | |
| MLLM2 | 22 39 | 18 37 | 18 41 | |
| OLDM2 | 18 31 | 13 28 | 12 35 | |
| OXNM2 | 23 42 | 22 40 | 23 48 | |
| PRAM2 | 22 49 | 22 45 | 18 45 | |

Western Pacific MOS Guidance




| NSTU | GFS MOS GUIDANCE | | | | | | | | | | | | | | | | | | | | | | | | |
|------|------------------|-------|----|----|----|----|----------|----|----|----|----|----|----|----|----|----|----|------|----|----|----|--|--|------|----|
| | 11/07/2008 | | | | | | 1200 UTC | | | | | | | | | | | | | | | | | | |
| DT | /NOV | 7/NOV | 8 | | | | | | | | | | | | | | | /NOV | 9 | | | | | /NOV | 10 |
| HR | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 06 | 12 | | | | |
| TMP | 84 | 85 | 85 | 85 | 82 | 82 | 81 | 79 | 80 | 83 | 84 | 83 | 81 | 81 | 80 | 79 | 81 | 84 | 86 | 82 | 80 | | | | |
| DPT | 77 | 77 | 78 | 77 | 76 | 77 | 76 | 75 | 77 | 78 | 77 | 77 | 76 | 77 | 76 | 75 | 77 | 78 | 77 | 77 | 76 | | | | |
| WDR | 08 | 08 | 08 | 09 | 08 | 07 | 05 | 04 | 06 | 07 | 08 | 07 | 05 | 02 | 35 | 01 | 02 | 07 | 07 | 08 | 10 | | | | |
| WSP | 17 | 17 | 15 | 13 | 11 | 08 | 07 | 07 | 07 | 08 | 09 | 08 | 07 | 05 | 04 | 04 | 04 | 08 | 09 | 06 | 06 | | | | |
| P06 | | 36 | | 37 | | 47 | | 46 | | 50 | | 43 | | 25 | | 35 | | 43 | | 30 | 31 | | | | |
| P12 | | | | 60 | | | | 66 | | | | | 60 | | | 59 | | | | | 47 | | | | |

Application of Linear Regression to MOS Development

MOS LINEAR REGRESSION

JANUARY 1 - JANUARY 30, 1994 0000 UTC


KCMH



MOS LINEAR REGRESSION


JANUARY 1 - JANUARY 30, 1994 0000 UTC

KCMH



REDUCTION OF VARIANCE


A measure of the “goodness” of fit and
Predictor / Predictand correlation



MOS LINEAR REGRESSION

JANUARY 1 - JANUARY 30, 1994 0000 UTC


KUIL



MOS LINEAR REGRESSION

DECEMBER 1 1993 - MARCH 5 1994 0000 UTC


KCMH



MOS LINEAR REGRESSION

DECEMBER 1 1993 - MARCH 5 1994 0000 UTC


KCMH



MOS LINEAR REGRESSION

DECEMBER 1 1993 - MARCH 5 1994 0000 UTC


KCMH



MOS LINEAR REGRESSION

DECEMBER 1 1993 - MARCH 5 1994 0000 UTC

KCMH



EXAMPLE REGRESSION EQUATIONS

$$Y = a + bX$$

CMH MAX TEMPERATURE EQUATION

$$\text{MAX T} = -352 + (0.3 \times 850 - 1000 \text{ mb THICKNESS})$$


CMH PROBABILITY OF PRECIPITATION EQUATION

$$\begin{aligned} \text{POP} = & -0.234 + (0.007 \times \text{MEAN RH}) \\ & + (0.478 \times \text{BINARY MEAN RH CUTOFF AT 70\%})^* \end{aligned}$$

* (IF MRH \geq 70% BINARY MRH = 1; else BINARY MRH = 0)

If the predictand is **BINARY**,
MOS regression equations produce
estimates of event **PROBABILITIES**...

KCMH



Making a PROBABILISTIC statement...

PEANUTS CHARLES M. SCHULZ



Quantifies the uncertainty !

DEFINITION of PROBABILITY

(Wilks, 2006)

- The degree of belief, or *quantified judgment*, about the occurrence of an uncertain event.

OR

- The long-term relative frequency of an event.

PROBABILITY FORECASTS

Some things to keep in mind

Assessment of probability is *EXTREMELY* dependent upon how predictand “event” is defined:



- Time period of consideration
- Area of occurrence
- Dependent upon another event?

MOS forecasts can be:

- POINT PROBABILITIES
- AREAL PROBABILITIES
- CONDITIONAL PROBABILITIES

AREAL PROBABILITIES

3H Eta MOS thunderstorm probability forecasts
valid 0000 UTC 8/27/2002 (21-24h proj)



What if these were 6-h forecasts?

40-km gridbox
10% contour interval


20-km gridbox
10% contour interval

PROPERTIES OF MOS PROBABILITY FORECASTS

- **Unbiased**
Average forecast probability equals long-term relative frequency of event
- **Reliable**
Conditionally or “Piecewise” unbiased over entire range of forecast probabilities
- **Reflect predictability of event**
Range narrows and approaches event RF as NWP model skill declines
 - extreme forecast projection
 - rare events

Reliable Probabilities...

Even for rare events



Designing an Operational MOS System:


Putting theory into practice...

DEVELOPMENTAL CONSIDERATIONS

MOS in the real world

- Selection (and QC!) of Suitable Observational Datasets
ASOS? Remote sensor? Which mesonet?


Suitable observations?



Appropriate Sensor?


Good siting?

Real or Memorex?



MOS Snowfall Guidance


Uses Observations from Cooperative Observer Network




36-hr forecast

12Z 12/05/03 – 12Z 12/06/03

Storm #2 Totals Dec 5-6, 2003



Map by: Chris Strong



Verification

DEVELOPMENTAL CONSIDERATIONS

MOS in the real world

- **Selection (and QC!) of Suitable Observational Datasets**
ASOS? Remote sensor? Which mesonet?
- **Predictand Definition**
Must be precise !!

PREDICTAND DEFINITION

Max/Min and PoP

Daytime Maximum Temperature

“Daytime” is 0700 AM - 0700 PM LST *

Nighttime Minimum Temperature

“Nighttime” is 0700 PM - 0800 AM LST *

* CONUS – differs in AK

Probability of Precipitation

Precipitation occurrence is accumulation of ≥ 0.01 inches of liquid-equivalent at a gauge location within a specified period


PREDICTAND DEFINITION

GFSX 12-h Average Cloud Amount


- Determined from 13 consecutive hourly ASOS observations, satellite augmented
- Assign value to each METAR report:
CLR; FEW; SCT; BKN; OVC
0 ; 0.15; 0.38; 0.69; 1
- Take weighted average of above
- Categorize:
CL < .3125 ≤ PC ≤ .6875 < OV

Creating a Gridded Predictand

Lightning strikes are summed over the “appropriate” time period and assigned to the center of “appropriate” grid boxes



A thunderstorm is deemed to have occurred when one or more lightning strikes are observed within a given gridbox:



= thunderstorm



= no thunderstorm

DEVELOPMENTAL CONSIDERATIONS

MOS in the real world

- **Selection (and QC!) of Suitable Observational Datasets**
ASOS? Remote sensor? Which mesonet?
- **Predictand Definition**
Must be precise !!
- **Choice of Predictors**
“Appropriate” formulation
Binary or other transform?

“APPROPRIATE” PREDICTORS

- DESCRIBE PHYSICAL PROCESSES ASSOCIATED WITH OCCURRENCE OF PREDICTAND

i.e. for POP:

PRECIPITABLE WATER
VERTICAL VELOCITY
MOISTURE DIVERGENCE
MODEL PRECIPITATION

~~1000-500 MB THK
TROPopause HGT~~

- “MIMIC” FORECASTER THOUGHT PROCESS
(VERTICAL VELOCITY) X (MEAN RH)

POINT BINARY PREDICTOR

24-H MEAN RH

CUTOFF = 70%

INTERPOLATE ; STATION RH \geq 70% , BINARY = 1
BINARY = 0 OTHERWISE

96

86

89

94

87

73

76

90

(71%) ● KCMH

76

60

69

92

64

54

68

93

RH \geq 70% ; BINARY AT KCMH = 1

GRID BINARY PREDICTOR

24 H MEAN RH CUTOFF = 70%

WHERE $RH \geq 70\%$; GRIDPOINT = 1 ; INTERPOLATE

1 1 1 1 1

1 1 1 1 1

1 0 0 1


0 0 0 1

(.21) • KCMH

$0 \leq \text{VALUE AT KCMH} \leq 1$

Logit Transform Example

KPIA (Peoria, IL) 0000 UTC ; 18-h projection



DEVELOPMENTAL CONSIDERATIONS

(cont.)

- **Terms in Equations; Selection Criteria**

“REAL” REGRESSION EQUATIONS

MOS regression equations are MULTIVARIATE , of form:

$$Y = a_0 + a_1 X_1 + a_2 X_2 + \dots + a_N X_N$$

Where,

the "a's" represent COEFFICIENTS


the "X's" represent PREDICTOR variables

The maximum number of terms, N , can be **QUITE** large:

For GFS QPF, $N = 15$ For GFS VIS, $N = 20$

The **FORWARD SELECTION** procedure determines the predictors and the order in which they appear.

FORWARD SELECTION

- METHOD OF PREDICTOR SELECTION ACCORDING TO CORRELATION WITH PREDICTAND
 - “BEST” OR STATISTICALLY MOST IMPORTANT PREDICTORS CHOSEN FIRST
-
- **FIRST** predictor selected accounts for greatest reduction of variance (RV)
 - Subsequent predictors chosen that give greatest RV in conjunction with predictors already selected
 -  selection when desired maximum number of terms is reached or new predictors provide less than a user-specified minimum RV

DEVELOPMENTAL CONSIDERATIONS


(cont.)

- **Terms in Equations; Selection Criteria**
- **Dependent Data**
 - Sample Size, Stability, Representativeness**
 - AVOID OVERFIT !!**
 - Stratification - Seasons**
 - Pooling – Regions**

MOS LINEAR REGRESSION


OCTOBER 1 1993 - MARCH 31 1994 0000 UTC

KUIL



GFS MOS Cool Season PoP/QPF Regions

With GFS MOS forecast sites (1720) + PRISM



DEVELOPMENTAL CONSIDERATIONS


(cont.)

- **Terms in Equations; Selection Criteria**
- **Dependent Data**
 - Sample Size, Stability, Representativeness**
 - AVOID OVERFIT !!**
 - Stratification - Seasons**
 - Pooling – Regions**
- **Categorical Forecasts?**

MOS BEST CATEGORY SELECTION

KDCA 12-Hour QPF Probabilities

48-Hour Projection valid 1200 UTC 10/31/93




How well do we do?

MOS Verification


Temperature Verification - 0000 UTC

GFS MOS vs. GFS DMO (4/2004 - 5/2006)




Temperature Verification - 0000 UTC

Warm Season: April – September, 2010




Temperature Bias - 0000 UTC

Warm Season: April – September, 2010




Temperature Bias - 0000 UTC

10/06; 01/07; 03/08



PoP Verification - 0000 UTC


10/06; 01/07; 03/08




GFSX 12-h Forecast Skill - 0000 UTC

Max Temperatures and PoP

% Improvement over Climate
Cool Season 1997 - 2003




Max T




PoP

45-yr Max Temperature Verification

Guidance / WFO; Cool Season 1966 - 2010



Dealing with NWP model changes



Mitigating the effects on development

To help reduce the impact of model changes and small sample size, we rely upon...

1. Improved model realism

better model = better statistical system

2. Coarse, consistent archive grid

smoothing of fine-scale detail

constant mesh length for grid-sensitive calculations

3. Enlarged geographic regions


larger data pools help to stabilize equations

4. Use of “robust” predictor variables

fewer boundary layer variables

variables likely immune to known model changes;

(e.g. combinations of state variables only)




Responding to NWP Model Changes

- **Parallel evaluation**


Run MOS...new vs. old NWP model

Assess impacts on MOS skill



Responding to NWP Model Changes

Eta MOS PoP: Eta vs. NMM output



Responding to NWP Model Changes


- **Parallel evaluation**

 - Run MOS...new vs. old NWP model
 - Assess impacts on MOS skill

- **Do nothing?**






 - OK if impacts are minimal

 - But, often they aren't! (GFS wind / temps)




2009 - 2011 GFS MOS Wind Bias

GFS MOS Annual Mean Wind Speed Biases For 24 Hrs Projection & 00Z Cycle (unit: KT)



Wind Speed Bias for KABQ July - Sept. 2010 (00Z Cycle)



Responding to NWP Model Changes


- **Parallel evaluation**

 - Run MOS...new vs. old NWP model
 - Assess impacts on MOS skill

- **Do nothing?**

 - OK if impacts are minimal

 - But, often they aren't! (GFS wind / temps)



- **OK, now what?**

 - Model changes may be recent


 - i.e. limited sample available from new model version

 - Error characteristics significantly different

 - Undesirable effects on MOS performance


Responding to NWP Model Changes

- Bias Correction for MOS?



Daily Bias Correction

based on past N (7, 10, 20 or 30)- day forecast errors




F = Forecasts ; O = Observations
N = Days in training sample
(typically, N = 7, 10, 20, or 30)

Daily biases can be treated equally or weighted to favor most recent days, etc.


Raw / Corrected GFS MOS Wind MAE

KABQ – 00UTC, 96-h Projection



Raw / Corrected GFS MOS Temp MAE

Southwest U.S. – 00UTC, 48-h Projection




Responding to NWP Model Changes

- **Bias Correction for MOS?**

Apply to Temps? Winds?

Run continuously in background?

Satisfactory in rapidly-varying conditions?



- **Redevelop?**


Short sample from new model or “mixed”?

Full System, selected elements?

Biggest impacts on single-station equations (Temp, Wind)

NAM / Eta MOS Dewpoint Comparison

Jul 15-31, 2006 and May 1-15, 2007



Using even just a little data from the new NWP model version can be helpful!

Eta MOS Replacement

December 9, 2008

“Classic” Eta MOS

| KORD | ETA MOS GUIDANCE | | | | | | | | | | | | 9/27/2007 1200 UTC | | | | | | | | |
|------|------------------|----|----|----|----|----|----------|----|----|----|----|----|--------------------|----|----|----|----|----|---------|----|----|
| DT | /SEPT 27/SEPT 28 | | | | | | /SEPT 29 | | | | | | /SEPT 30 | | | | | | | | |
| HR | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 06 | 12 |
| N/X | | | | | | 50 | | | | | 72 | | | | 50 | | | | 78 | 57 | |
| TMP | 67 | 69 | 64 | 60 | 56 | 53 | 52 | 64 | 70 | 70 | 65 | 58 | 55 | 53 | 53 | 65 | 74 | 77 | 71 | 61 | 58 |
| DPT | 54 | 52 | 51 | 50 | 48 | 46 | 46 | 48 | 45 | 44 | 44 | 47 | 47 | 47 | 48 | 50 | 49 | 49 | 51 | 55 | 53 |
| CLD | OV | BK | SC | SC | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | FW |
| WDR | 23 | 27 | 30 | 30 | 29 | 28 | 30 | 32 | 33 | 34 | 06 | 11 | 16 | 17 | 17 | 18 | 18 | 19 | 17 | 19 | 20 |
| WSP | 09 | 10 | 08 | 06 | 05 | 05 | 04 | 07 | 08 | 07 | 06 | 02 | 02 | 02 | 04 | 08 | 09 | 10 | 08 | 08 | 08 |
| P06 | | 19 | | 3 | | 6 | | 1 | | 0 | | 0 | | 1 | | 3 | | 3 | 8 | 10 | |
| P12 | | | | | | 6 | | | | 1 | | | | 1 | | | | 6 | | 12 | |
| Q06 | | 0 | | 0 | | 0 | | 0 | | 0 | | | | 0 | | 0 | | 0 | 0 | 0 | |
| Q12 | | | | | | 0 | | | | 0 | | | | 0 | | | | 0 | | 0 | |
| T06 | | 1/ | 0 | 9/ | 7 | 0/ | 0 | 0/ | 7 | 0/ | 0 | 0/ | 1 | 0/ | 0 | 0/ | 8 | 2/ | 0999/99 | | |
| T12 | | | | 9/ | 7 | | 0/ | 7 | | 0/ | 1 | | 0/ | 1 | | 0/ | 8 | | 999/99 | | |
| SNW | | | | | | 0 | | | | | | | | 0 | | | | | | 0 | |
| CIG | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| VIS | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| OBV | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |

- All Eta model input
- All elements used eta-based equations

Hybrid “NAM MOS”

| KORD | NMM MOS GUIDANCE | | | | | | | | | | | | 9/27/2007 1200 UTC | | | | | | | | | |
|------|------------------|----|----|----|----|----|----------|----|----|----|----|----|--------------------|----|----|----|----|----|--------|----|---------|----|
| DT | /SEPT 27/SEPT 28 | | | | | | /SEPT 29 | | | | | | /SEPT 30 | | | | | | | | | |
| HR | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 03 | 06 | 09 | 12 | 15 | 18 | 21 | 00 | 06 | 12 | |
| N/X | | | | | | 51 | | | | | 74 | | | | 49 | | | | 76 | 58 | | |
| TMP | 68 | 65 | 62 | 60 | 57 | 52 | 52 | 63 | 71 | 73 | 69 | 59 | 53 | 51 | 50 | 63 | 72 | 76 | 73 | 60 | 58 | |
| DPT | 54 | 53 | 53 | 51 | 49 | 48 | 48 | 48 | 47 | 45 | 45 | 47 | 47 | 47 | 48 | 50 | 50 | 50 | 51 | 52 | 55 | |
| CLD | OV | OV | SC | FW | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | CL | FW | FW | |
| WDR | 23 | 27 | 30 | 30 | 29 | 28 | 30 | 32 | 33 | 34 | 04 | 10 | 16 | 16 | 17 | 18 | 18 | 19 | 17 | 19 | 20 | |
| WSP | 08 | 10 | 08 | 06 | 05 | 05 | 05 | 07 | 08 | 07 | 05 | 03 | 02 | 03 | 04 | 08 | 09 | 10 | 08 | 08 | 08 | |
| P06 | | | 21 | | 3 | | | 2 | | 1 | | 1 | | 0 | | 0 | | 3 | | 5 | 10 | 12 |
| P12 | | | | | | 3 | | | | | | 1 | | | | 0 | | | | 7 | | 16 |
| Q06 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | | 0 | 0 | 0 |
| Q12 | | | | | | 0 | | | | | | 0 | | | | 0 | | | | 0 | | 0 |
| T06 | | 1/ | 0 | 6/ | 1 | 0/ | 0 | 0/ | 3 | 0/ | 0 | 0/ | 0 | 0/ | 0 | 0/ | 0 | 0/ | 1 | 4/ | 0999/99 | |
| T12 | | | | 6/ | 1 | | | 0/ | 3 | | 0/ | 0 | | 0/ | 0 | 0/ | 1 | | 999/99 | | | |
| SNW | | | | | | 0 | | | | | | | | 0 | | | | 0 | | | | 0 |
| CIG | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 | 8 |
| VIS | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 | 7 |
| OBV | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N | N |

- All NMM model input
- Redeveloped elements use new NMM-based equations
- Other elements use older Eta-based equations applied to NMM


Responding to NWP Model Changes

- **Bias Correction for MOS?**

Apply to Temps? Winds?

Run continuously in background?

Satisfactory in rapidly-varying conditions?



- **Redevelop?**

Short sample from new model or “mixed”?

Full System, selected elements?

Biggest impacts on single-station equations (Temp, Wind)

- **Reforecasts?**

1-2 year sample probably sufficient for T, Wind

Rare elements need longer or “mixed” sample?

Requires additional supercomputer resources

MOS: Today and Beyond

The Future of MOS

“Traditional” Station-oriented Products

- **GFS / GFSX MOS:**

 - Update GFSX Sky Cover equations

 - (Completes 1200 UTC text message)

 - Expand GFSX to Day 10 for some elements

 - Update climate normals (1981-2010 NCDC)

 - Bias-corrected T, Td, Max/Min, windspeed

- **NAM MOS (Eta MOS replacement):**

 - Add precipitation type suite (TYP, POZ, POS)

 - Add 0600 and 1800 UTC cycles?

 - Update remaining eta-based elements


 - Update temperature suite with NMM-b data

The Future of MOS

“Traditional” Station-oriented Products (contd.)


- **Western Pacific MOS:**
 - Add new elements (Sky Cover, CIG)
- **General:**
 - Evaluate impacts of NWP model changes
 - Periodic addition of new CONUS sites
 - Gradual phaseout of station-oriented graphics

GFS MOS 24-hr Conditional Probability of Precipitation ≥ 0.25 "



<http://www.nws.noaa.gov/mdl/synop>


End of an era?



WANTED! High-resolution, gridded guidance for NDFD

Gridded MOS

GFS-based CONUS-wide @ 5km



Max / Min

PoP

Temp / Td

RH

Tstm

Winds

QPF

Snowfall


Gusts

Sky Cover

[http://www.weather.gov/mdl/synop/
gridded/sectors/index.php](http://www.weather.gov/mdl/synop/gridded/sectors/index.php)


Wait...Stop the Presses!!

2.5-km CONUS GMOS - "live" on Feb. 27, 2012



Alaska Gridded MOS


AK GMOS: GFS-based, 3-km grid



**All elements complete
January, 2010**

Hawaii Gridded MOS

Hawaii GMOS: GFS-based, 2.5-km grid



Max / Min

PoP

Temp / Td

RH

Winds

Gusts

2.5-km grid

Implemented
November, 2010





The Future of MOS





“Enhanced-Resolution” Gridded MOS Systems

- “MOS at any point” (e.g. GMOS)
 - Support NWS digital forecast database
 - 2.5 km - 5 km resolution
 - Equations valid *away* from observing sites
 - Emphasis on high-density surface networks
 - Use high-resolution geophysical data

Surface observation systems used in GMOS





- METAR
- Buoys/C-MAN
- Mesonet (RAWS/SNOTEL/Other)
- NOAA cooperative observer network
- RFC-supplied sites





- GFS/NAM
- ▲ Marine
- Cooperative

MOS Stations




**Approx.
11,000 sites!**


- GFS/NAM
- ▲ Marine
- Cooperative
- Mesonet
- RFC

MOS Stations


Gridded MOS – Central CA



Geophysical Datasets




5-km Terrain




5-km Land Cover

Gridded MOS Concept - Step 1

“Blending” first guess and high-density station forecasts



First guess field from
Generalized Operator Equation
or other source



First guess + guidance
at all available sites


Developing the “First Guess” Field

Some options


- **Generalized operator equation (GOE)**
 - Pool observations regionally
 - Develop equations for all elements, projections
 - Apply equations at all grid points within region
- **Use average field value at all stations**
- **Use other user-specified constant**
- **Use NWP model forecast**

Gridded MOS Concept - Step 2

Add further detail to analysis with high-resolution geophysical data and “smart” interpolation



First guess + guidance
at all available sites



First guess + station forecasts +
terrain

GMOS Analysis

Basic Methodology (Glahn, et al. 2009, WaF)

- **Method of successive corrections (“BCDG”)**
Bergthorssen and Doos (1955); Cressman (1959);
Glahn (1985, LAMP vertical adjustment)
- **Elevation (“lapse rate”) adjustment**
Inferred from forecasts at different elevations
Calculations done “on the fly” from station data
Can vary by specific element, synoptic situation
- **Land/water gridpoints treated differently**

GMOS Analysis

Other Features

- **Special, terrain-following smoother**
- **ROI can be adjusted to account for variations in density of observed data**
- **Nudging can be performed to help preserve nearby station data**
- **Parameters can be adjusted individually for each weather element**


GMOS Analysis

Some Issues


- **Not optimized for all weather elements and synoptic situations**
Need situation specific, dynamic models?
- **May not capture localized variations in vertical structure**
Vertical adjustment uses several station “neighbors”
- **May have problems in data-sparse regions over flat terrain**
Defaults to pure Cressman analysis with small ROI
Can result in some “bulls-eye” features

NDGD vs. NDFD

Which is "better"?




NDGD Max T




NDFD Max T

NDGD vs. NDFD

Which is “better”?



NDGD RH




NDFD RH

Fewer obs available to analysis = less detail in GMOS

Forecasters adding detail: Which is “better”? More accurate?

AK GMOS Temps & Observing Sites

Even fewer obs available – Yikes!




The Future of MOS

“Enhanced-Resolution”, Gridded MOS Systems

- “MOS at any point” (e.g. GMOS)
 - Support NWS digital forecast database
 - 2.5 km - 5 km resolution**
 - Equations valid **away** from observing sites
 - Emphasis on high-density surface networks
 - Use high-resolution geophysical data
- “True” gridded MOS
 - Observations and forecasts valid on fine grid
 - Use remotely-sensed predictand data
 - e.g. WSR-88D QPE, Satellite clouds, NLDN**

Remotely-sensed precipitation data



- ≤ 12.00
- ≤ 10.00
- ≤ 8.00
- ≤ 6.00
- ≤ 5.00
- ≤ 4.00
- ≤ 3.00
- ≤ 2.00
- ≤ 1.50
- ≤ 1.00
- ≤ 0.75
- ≤ 0.50
- ≤ 0.25
- ≤ 0.10
- ≤ 0.01


The Future of MOS

Gridded MOS: Where do we go from here?


- **Additions to current CONUS GMOS system**
 - Present weather grid**
 - NAM-based companion system (short-range)**
 - Probabilistic and/or ensemble-based products**

Sample Forecast as Quantile Function (CDF)

(72-h Temp KBWI 12/14/2004)



Ensemble MOS [5%, 95%] Quantile Temps.



The Future of MOS

Gridded MOS: Where do we go from here?

- Additions to current CONUS GMOS system
 - “Predominant” weather grid
 - NAM-based companion system (short-range)
 - Probabilistic and/or ensemble-based products
- Increase CONUS resolution from 5-km to 2.5-km
 - NCEP jobstream Feb. 2012; awaits comms upgrade
- Update land / water mask based on WFO input
- Improve GMOS interpolation procedures

The Future of MOS

Gridded MOS: Where do we go from here?

- Increase utilization of mesonet data
Investigate MADIS archive (NCO/TOC/ESRL)
~20,000 additional sites?
- Incorporate remotely-sensed data where possible
SCP augmented clouds (already in use)
WSR-88D QPF: **March 13, 2012**
NSSL MRMS (Multi-radar, Multi-sensor) dataset?
- Expand GMOS for AK; add other OCONUS areas
AK: Increase grid extent; improve marine winds
Hawaii: add QPF, Sky Cover
Puerto Rico

REFERENCES...the “classics”

Wilks, D.: **Statistical Methods in the Atmospheric Sciences**, 2nd Ed., Chap. 6, p. 179 - 254.

Draper, N.R., and H. Smith: **Applied Regression Analysis**, Chap. 6, p. 307 - 308.

Glahn, H.R., and D. Lowry, 1972: The use of model output statistics in objective weather forecasting, JAM, 11, 1203 - 1211.

Carter, G.M., et al., 1989: Statistical forecasts based on the NMC's NWP System, Wea. & Forecasting, 4, 401 - 412.

REFERENCES (GMOS)

Glahn, H.R., et al., 2009: The Gridding of MOS.,
Wea. & Forecasting, 24, 520 – 529.