

GIS Applications in NWS Operations & Lake Effect Snow Forecasting

ATM 362 – Forecasting & NWS Operations Course

Joe Villani

Meteorologist - WFO Albany NY

March 2, 2020





Outline


- Introduction to GIS
- Elevation mapping
- Flash Flood Potential Index
- Rainfall/Snowfall analysis maps
- NWS zone-based verification
- NWS forecast error maps
- Snowfall composites stratified by flow regime
- Hi-res model verification
- Automation → “GAZPACHO” program
- Forecasting Lake Effect Snow (Albany forecast area)



Introduction to Geographic Information Systems

- What is GIS?


From ESRI.com: “A geographic information system (GIS) is a framework for gathering, managing, and analyzing data. Rooted in the science of geography, GIS integrates many types of data. It analyzes spatial location and organizes layers of information into visualizations using maps and 3D scenes. With this unique capability, GIS reveals deeper insights into data, such as patterns, relationships, and situations—helping users make smarter decisions.”





Introduction to GIS


- ArcMap is the main ArcGIS software used at NWS Albany (contains ArcCatalog for organizing GIS files)
- Powerful mapping application: creation of shape files → **rasters** (digital dataset made up of grid points)
- Crucial aspect of Rasters → can be mathematically manipulated





Local Applications: Elevation Maps


- Uses USGS Digital Elevation Model (DEM) data 30 meter resolution
- Download in tiles, then mosaic using ArcMap




DEM Mosaic in ArcMap

Local Applications: Elevation Maps

- Delineate elevation thresholds in 500 ft increments for elevation-specific maps




> 1000 ft



> 2000 ft


Local Applications: Elevation Maps

- Create contoured elevation map with 500 ft range bins
- Select different color for each range bin
- Useful for visualizing key topographic features in the Albany forecast area




Flash Flood Potential Index (FFPI)

- The FFPI incorporates physiographic characteristics of an individual drainage basin to determine its hydrologic response
- For Flash Flooding, hydrologic response influenced by four main factors:
 1. Soil type
 2. Terrain slope
 3. Vegetation and forest canopy
 4. Land use, especially urbanization



Flash Flood Potential Index (FFPI)

- High resolution maps obtained and converted to raster datasets using GIS software (ArcMap)
- Data processed over the domain of interest (ALY forecast area)
- Use ArcMap tools to resample, reclassify and combine the data
- Result is a numerical index of flash flood potential specific to the ALY forecast area







Lowest Flash
Flood potential

Highest Flash
Flood potential




Flash Flood Potential Index (FFPI)

- Combine slope, land use, soil type and forest density using the following formula:
- $\text{FFPI} = (1.5 * \text{Slope} + \text{LU} + \text{FD} + \text{Soil}) / N$
- Average the index over each basin



Flash Flood Potential Index (FFPI)

- Helps identify areas of greater/lesser Flash Flood Potential
- Areas of steep slope, agricultural use, and urban areas have higher FF potential
- Areas of dense forest, lakes/ponds, and less slope have lower FF potential




Rainfall/Snowfall Analysis Maps

- Create interpolated rainfall/snow maps from observations

- Compile snow/rain reports from Public Information Statement
- Generate GIS shape file of snowfall reports from spreadsheet using ArcCatalog
- Import into ArcMap

A	B	C	D	E	F
Site name	Snowfall	Source	LAT	LON	LOC
Bakersville	6.2	Co-Op Observer	41.85833	-73.0103	ALY
New Hartford	6.2	CoCoRaHS	41.8417	-73.0092	ALY
Canaan	6	WeatherNet6	42.01215	-73.3505	ALY
Norfolk	6	Co-Op Observer	41.9725	-73.2208	ALY
Winsted	6	CoCoRaHS	42.002	-73.0825	ALY
Litchfield	4.9	CoCoRaHS	41.77445	-73.169	ALY
Woodbury Center	3.5	Amateur Radio	41.5558	-73.2261	ALY
New Milford	3	CoCoRaHS	41.51472	-73.436	ALY
New Milford	3	Amateur Radio	41.5885	-73.4068	ALY
Roxbury	3	Amateur Radio	41.5539	-73.305	ALY
New Milford	2.6	CoCoRaHS	41.59392	-73.4515	ALY
Woodbury Center	2	Amateur Radio	41.5466	-73.207	ALY
Savoy	20.3	WeatherNet6	42.59784	-73.0436	ALY
Pittsfield	7.2	Trained Spotter	42.4518	-73.2609	ALY
Lenoxdale	7	Social Media	42.337	-73.2456	ALY
Pittsfield	6.5	CoCoRaHS	42.47878	-73.2738	ALY
Housatonic	6	CoCoRaHS	42.2379	-73.3503	ALY
Housatonic	6	CoCoRaHS	42.28983	-73.3195	ALY
Sandisfield	4.5	Social Media	42.09872	-73.1377	ALY






Rainfall/Snowfall Analysis Maps



- Run an Inverse Distance Weighting (IDW) function in ArcMap
- IDW takes into account distance between points & interpolates
- Creates a Raster (digital) dataset and contoured rainfall or snowfall map




Snowfall Example



Rainfall/Snowfall Analysis Maps

- Run an Inverse Distance Weighting (IDW) function in ArcMap
- IDW takes into account distance between points & interpolates
- Creates a Raster (digital) dataset and contoured rainfall or snowfall map



Rainfall Example

Zone-based Snowfall Verification

- NWS Verification computed from mean snowfall in each forecast zone
- Run Zonal Statistics in ArcMap (including mean) on the observed snowfall analysis Raster





Image source: <http://help.arcgis.com>

Zone-based Snowfall Verification

- Output via map, with more specific data in spreadsheet format
- **Mean** snowfall for each zone used for NWS snowfall verification




A	B	C	D	E	F
NAME	MIN	MAX	RANGE	MEAN	STD
NY083:Southeast Warren	12.1	23.4	11.3	21	1.4
NY050:Southern Saratoga	13	21.1	8.1	18	1.3
NY057:Delaware	7.9	17	9.1	12.3	1.8
NY060:Western Columbia	5.6	12	6.4	8.9	1.3
NY038:Southern Herkimer	10.3	20.4	10.1	15.6	2.7
NH007:Sullivan	7.4	13.4	5.9	10.6	1.5
NY052:Eastern Albany	10.4	17.2	6.8	13.9	0.9
NY032:Northern Herkimer	11	19.3	8.3	15.4	2.5
NY046:Otsego	8.2	19.1	10.9	13.5	2.2
NY029:Southeastern St. Lawrence	10	13.9	3.9	12.2	0.7
NY042:Northern Warren	13	23.4	10.5	19.5	2.1
VT015:Eastern Windham	9.4	15.4	6.0	12.7	1


Creation of Forecast Error Maps

- Methodology:
Create gridded observed snowfall map after an event, then subtract from the forecast snowfall (obtained via NWS **National Digital Forecast Database**)


NDFD Forecast Snow



Minus



Observed Snow



Difference


Monthly Rainfall/Snowfall Maps

Rainfall November 2019

Snowfall December 2019

National Weather Service Albany New York
Monthly Liquid Equivalent Precip. November 2019

Analysis Data Source: Regional Observations




Rainfall Analysis 0" - 50"

- 0.00"
- 0.01" - 0.09"
- 0.10" - 0.24"
- 0.25" - 0.49"
- 0.50" - 0.99"
- 1.00" - 1.49"
- 1.50" - 1.99"
- 2.00" - 2.99"
- 3.00" - 3.99"
- 4.00" - 5.99"
- 6.00" - 7.99"
- 8.00" - 9.99"
- 10.00" - 14.99"
- 15.00" - 19.99"
- 20.00" - 29.99"
- 30.00" - 50.00"

Created:
12/22/2019 08:57AM

National Weather Service Albany New York
Monthly Snowfall December 2019

Analysis Data Source: Regional Observations



Snowfall Analysis 0" - 100"

- 0.0"
- < 1.0"
- 1.0" - 2.0"
- 2.0" - 3.0"
- 3.0" - 4.0"
- 4.0" - 6.0"
- 6.0" - 8.0"
- 8.0" - 12.0"
- 12.0" - 18.0"
- 18.0" - 24.0"
- 24.0" - 30.0"
- 30.0" - 36.0"
- 36.0" - 50.0"
- 50.0" - 100.0"


Created:
01/10/2020 05:55AM

- Same method used for event precipitation analysis maps, but instead using monthly rainfall/snowfall reports (based on local reports from observers)

Seasonal Snowfall Map

- Methodology:
Compile (SUM)
monthly
observed
snowfall maps
from October
through April
using ArcMap
- Color scale &
range bins must
be altered to
account for
large snow
totals

2018-19 Seasonal Snowfall






Climatology Snowfall Map

1981-2010 Seasonal Snowfall Climatology

- Use 30-year climate normals from Cooperative Weather Observers to create a snowfall analysis





Snowfall Composites Stratified by 925mb Flow Regime



- Identify patterns of snowfall distribution based on low level flow regime
- Use Albany, NY sounding data to determine wind at 925 mb
- Sounding time (0000 UTC or 1200 UTC) closest to midpoint of each event

KALB (72518) 30/00 4245 -7348 92m LI (500mb):							
WMO PARTS INCLUDED:		TTAA	TTBB	PPBB	TTCC	TTDD	PPDD
PRESS	TEMP	DWPT	WDIR	WSPD	HGHT	THTA	
MB	C	C	DEG	KNTS	M	K	
1010.0	17.4	12.7	0.0	0.0	92.0	289.7	
1000.0	16.2	11.2	35.0	3.0	179.0	289.4	
985.3	15.4	9.1	55.0	5.0	304.8	289.7	
965.0	14.2	6.2	92.1	5.6	481.3	290.3	
950.5	13.0	6.4	110.0	7.0	609.6	290.4	
925.0	11.0	6.7	130.0	11.0	837.0	290.6	
923.0	11.0	7.0	131.3	11.2	856.4	290.7	
916.6	10.8	6.6	135.0	12.0	914.4	291.1	
892.0	9.8	5.1	154.0	8.0	1140.6	292.3	
886.0	9.8	0.8	161.6	7.3	1196.9	292.9	
883.6	9.7	0.9	165.0	7.0	1219.2	293.0	
875.0	9.2	1.2	159.4	6.6	1300.4	293.3	
865.0	9.2	-5.8	152.1	6.3	1395.5	294.3	
850.0	8.8	-7.2	140.0	6.0	1540.0	295.4	
844.0	8.4	-7.6	136.4	4.2	1602.2	295.5	
829.0	9.0	-16.0	28.3	1.0	1751.8	297.7	
821.3	8.7	-16.7	345.0	3.0	1828.8	298.2	
792.0	7.6	-19.4	149.4	1.9	2126.8	300.1	
791.3	7.6	-19.4	150.0	2.0	2133.6	300.1	
783.0	7.0	-19.0	168.3	1.7	2219.7	300.4	
762.3	5.1	-8.8	220.0	2.0	2438.4	300.7	
759.0	4.8	-7.2	227.7	2.0	2473.7	300.7	
746.0	5.2	-3.8	254.3	2.4	2614.1	302.7	
734.2	4.2	-4.8	270.0	3.0	2743.2	303.0	
707.0	2.0	-7.0	180.0	3.0	3048.0	303.8	
700.0	1.4	-7.6	165.0	7.0	3128.0	304.0	

Example of Data Selection




Snowfall Composites Stratified by 925 mb Flow Regime



- Stratify by wind direction & speed at 925 mb:
 - 4 quadrants for wind direction
 - 3 categories for speed
- = 12 combined categories

Quadrants



Speed category	
1	0-19 kt
2	20-39 kt
3	40+ kt



Snowfall Composites Stratified by 925 mb Flow Regime




	A	B	C	D	E	F	G
1	925 mb	I-1	I-2	I-3	II-1	II-2	
2		20170331_0600-20170401_1800	20170123_1800-20170125_1800	20130307_1800-20130308_1800	20170212_1200-20170213_1800	20170117_1800-20170118_1800	
3		20170314_0000-20170315_1200	20160205_0600-20160205_1800		20170131_1200-20170201_1200	20161207_0000-20161207_1800	
4		20170107_0000-20170108_1200	20150126_1800-20150128_0600		20150208_0600-20150210_1200	20161205_0000-20161205_1800	
5		20160208_1200-20160209_1200	20141209_0600-20141211_1200		20140204_1800-20140206_0000	20150202_0000-20150203_0000	
6		20160117_1800-20160118_1200	20140312_1200-20140313_1800			20131214_1200-20131215_1800	
7		20140121_1800-20140122_1200	20140213_1200-20140214_1800				
8		20140102_0000-20140103_1800	20130318_1800-20170319_1800				
9		20130208_0600-20130209_1800					
10							
11							
12							
13							
14							
15	850 mb	I-1	I-2	I-3	II-1	II-2	
16		20170209_0600-20170210_0000	20150126_1800-20150128_0600	20140213_1200-20140214_1800	20170331_0600-20170401_1800	20170212_1200-20170213_1800	
17		20170107_0000-20170108_1200	20141209_0600-20141211_1200		20170314_0000-20170315_1200	20170123_1800-20170125_1800	
18		20140121_1800-20140122_1200	20130318_1800-20170319_1800		20140204_1800-20140206_0000	20160208_1200-20160209_1200	
19			20130307_1800-20130308_1800			20140102_0000-20140103_1800	
20						20131214_1200-20131215_1800	
21						20130208_0600-20130209_1800	

- Separate events into the 12 categories using a spreadsheet
- 81 total events from 2014-2019 winter seasons

Snowfall Composites Stratified by 925 mb Flow Regime


- Sum the snowfall analyses for events in each of the 12 categories using 925 mb wind at Albany



Snowfall Composites: Notable Regimes

Moderate SE Flow


2013-19 Snowfall Composite (in) - 925 mb Cat. ii-2 (10 events)
(90-180 degrees, 20-39 kt)



**Maximum snowfall in favored
upslope areas north and east of
Albany**

Moderate SW Flow

2013-19 Snowfall Composite (in) - 925 mb Cat. iii-2 (10 events)
(180-270 degrees, 20-39 kt)




**“Snow hole” in Capital District due
to down-sloping off Catskills**

Snowfall Composites: Notable Regimes


Moderate NE Flow

2013-19 Snowfall Composite (in) - 925 mb Cat. i-2 (8 events)
(0-90 degrees, 20-39 kt)



Moderate NW Flow

2013-19 Snowfall Composite (in) - 925 mb Cat. iv-2 (15 events)
(270-360 degrees, 20-39 kt)




**Snowfall dominated by
Nor'Easters**

**Maximum snowfall in favored
upslope areas of Adirondacks and
Taconics, Green Mountains &
Berkshires**

Snowfall Error Maps Stratified by Flow Regime


- Same methodology as snowfall composites, but summing difference maps for events in each of the 12 categories using 925 mb wind at Albany
- Used for assessing +/- forecast biases



Error Maps: Notable Regimes


Weak NE Flow

2013-19 Snowfall Bias (in) - 925 mb Cat. i-1 (10 events)
(0-90 degrees, 0-19 kt)



Moderate NW Flow

2013-19 Snowfall Bias (in) - 925 mb Cat. iv-2 (16 events)
(270-360 degrees, 20-39 kt)



Negative/positive snowfall forecast errors likely due to max QPF displacement error


**Under-forecast of upslope and lake effect snow
Over-forecast in “snow shadowing” areas**



HRRR and NAMNest Snowfall Verification




- NWS Albany verifies snowfall from high-resolution models such as the 3 km HRRR and NAMNest
- Use **positive snow depth change** from beginning to end of an event



HRRR

1-3 Dec 2019



NAMNest

1-3 Dec 2019




HRRR and NAMNest Snowfall Verification



- The change in snow depth more accurately depicts model snowfall (correspondence with EMC)
- Subtract observed snowfall from model positive snow depth change (using ArcMap) for each event to determine model forecast error


HRRR Pos SD Change



Minus



Observed Snow




Difference



HRRR and NAMNest Case Study


HRRR error

15-16 November 2018



NAMNest error

15-16 November 2018




- The positive SD change from both HRRR and NAMNest severely underestimated snowfall during the 14-15 Nov 2018 snowfall event
- Models likely forecasted SLR to be much too low across southern part of the area, where the heaviest snow fell




HRRR and NAMNest Case Study


HRRR Positive SD Change



NAMNest Positive SD Change



Observed Snowfall



15-16 November 2018




HRRR and NAMNest Snowfall Verification




HRRR composite bias (13 events)

HRRR Snowfal Bias 2018-19: Nov 15-16, Nov 19-20, Nov 26-28, Jan 3, Jan 9-10, Jan 19-20, Jan 27-28, Jan 29-30, Feb 12-13, Feb 27-28, Mar 3-4, Mar 10, Mar 22-23 (13 events)



NAMNest composite bias (13 events)

NAMNest Snowfal Bias 2018-19: Nov 15-16, Nov 19-20, Nov 26-28, Jan 3, Jan 9-10, Jan 19-20, Jan 27-28, Jan 29-30, Feb 12-13, Feb 27-28, Mar 3-4, Mar 10, Mar 22-23 (13 events)




- Sum difference maps (using ArcMap) to get seasonal composite snowfall bias for the HRRR and NAMNest
- Noticed a large negative (under-forecast bias). Seemed most extreme with early season events and lake effect

Automation using GIS Scripting: The “GAZPACHO” Program

- Not talking about cold soup here...
- **GAZPACHO** = **G**ridded **A**utomated **Z**onal **P**recipitation **A**nd **C**omplete **H**i-res **O**utput
- GAZPACHO is a program that was created to assist NWS Weather Forecast Offices with snowfall verification (rainfall added)



National Weather Service Albany New York
NDFD Snowfall Verification 1-3 December 2019
Analysis Data Source: Regional Observations





Automation using GIS Scripting: The “GAZPACHO” Program



- Created by Danny Gant (Morristown, TN), Vasil Koleci & Joe Villani (Albany, NY)
- Uses ArcGIS software and Python scripts
- GAZPACHO is run on a PC (with ArcGIS 10.7 installed) via a simple GUI

Required Fields

Choose Phenomena Type(s) To Verify Choose Phenomena Analysis Data Source(s)

Rainfall Snowfall IceAccum PNS/Other NOHRSC/AHPS Blend

Select or Enter Start Date Select or Enter End Date

February 2019

Sun	Mon	Tue	Wed	Thu	Fri	Sat
				01	02	
03	04	05	06	07	08	09
10	11	12	13	14	15	16
17	18	19	20	21	22	23
24	25	26	27	28		

YYYY MM DD YYYY MM DD

Select Start Time > 0000 None Select End Time > 0000

Archive Sources > Locate Data File xmACIS

Real-Time Sources > IRIS MRMS > Over Previous (N) Hours > 24

Optional Fields

Elevation Threshold (Feet) > Min Stats Only Max <

Incorporate Topography Enhancement Yes Only Elevations >

Only Zero Values From NOHRSC/AHPS Yes

Analysis Smoothing Factor None 2 4 8 16

States Counties Zones Points Cities Interstates Lakes Rivers

Enter Custom Headline

Align Fields From Custom Data File (Red=Required)

SiteName	Phenomena	Source	Latitude	Longitude	Location
None	None	None	None	None	None

Create Datafile

Archived Verification Data

Locate Archive File 1 Archive Type Locate Archive File 2

None


Close Download Model Data Create Maps





What is GAZPACHO


- Creates maps of:
- Observed precipitation (rain or snow)
- Zone-average rain/snow
- NWS forecast or model rain/snow (HRRR, NAMNest and NAM)
- Difference (or error) maps of forecast minus observed rain/snow (inches and %)
- Spreadsheet table of zone-average statistics





Case Study using GAZPACHO


- **Snowfall analysis** based on interpolated observations from 14-15 March 2017 Nor'easter (Pi Day Storm)



**Observed Snowfall
14-15 March 2017**

Case Study using GAZPACHO


- **Difference map** →
computed from
official NWS Albany
forecast minus
observed analysis
- Large under-forecast
error across areas
west of the Hudson
Valley
 - Mesoscale snow
band position &
magnitude errors



Forecast Error
14-15 March 2017

Case Study using GAZPACHO


- Snowfall  analysis based on interpolated observations from 02 March 2018
Nor'easter



Observed Snowfall
02 March 2018

Case Study using GAZPACHO


- **Difference map** →
computed from
official NWS Albany
forecast minus
observed analysis
- Large under-forecast
error in the Capital
District due to poor
model representation
of boundary layer
temps (esp. GFS)
- Heavy mesoscale
snow band just west
of Capital District



Forecast Error
02 March 2018

Forecasting Lake Effect Snow

- Instability Class
(derived from lake/air temp differences) used to assess potential strength of lake effect snow bands
- Three main categories:
 - 1) Conditional
 - 2) Moderate
 - 3) Extreme




Niziol Instability Class



Forecasting Lake Effect Snow


- From Nizioł, 1987
- **Conditional:** Heavy snowfall rates possible with trajectories along the longest fetch of the lake and 500 mb cyclonic vorticity advection (CVA)
- **Moderate:** Heavy snowfall rates can occur with moderately long trajectories & 500 mb CVA
- **Extreme:** Heavy snowfall rates possible with little dependence on fetch or CVA



BUFKIT – Instability category (must obtain observed lake temperature too)

Forecasting Lake Effect Snow

- Inversion heights also used to determine potential strength of lake effect snow bands
- Higher inversion heights result in a deeper mixed layer and taller convective plumes
- Inversion heights of 750 mb (~2.5 km) and higher can be associated with intense snow bands, assuming other favorable factors are present





BUFKIT – KRME NAM 26 December 2017



Forecasting Lake Effect Snow

Flow trajectory can be determined by using the mean wind in the mixed layer. Flow trajectory and fetch modulate the location and classification of snow bands as:

Single Band: wider in shape; typically more intense




Multi Band: narrower; typically weaker due to smaller fetch




Forecasting Lake Effect Snow

- Multi bands typically affect the southern Mohawk Valley, Schoharie Valley, and eastern Catskills with a 300° to 310° flow trajectory



- Single bands can affect the central and northern Mohawk Valley (even rarely the Capital District), and especially the western Adirondacks with a 260° to 280° flow trajectory



Limiting Factors



- Shallow inversion heights result in very weak snow showers/flurries
- If the directional wind shear is $> 30^\circ$ snow bands tend to become less organized or break up
- Diurnal mode especially early/late season, as bands tend to become cellular during the daylight hours


Forecast Challenges




- Inland extent of snow bands: Not always modulated by wind speed; can lead to over-forecasting significant events
- Model forecasts of QPF associated with LES can be highly inaccurate at times
- Snow to liquid ratios can range from 15:1 to as much as 30:1 (even rare occasions of $> 50:1$)

Favorable Env. far-reaching Inland Extent

- Type A
- MLC present (not shown)
- Strong 0-1 km shear; weak shear in 1-3 km layer
- Conditional lake-induced instability



(a)




(b)


(a) Example of a Type A sounding (0-hr NAM sounding from near KUCA) featuring strong 0–1-km bulk shear (weak in 1–3-km layer) and conditional instability. (b) Associated 0.5° reflectivity (dBZ) at time of sounding. Sounding and radar image from 1200 UTC 2 January 2012

Non-Favorable Env. far-reaching Inland Extent

- Type B
- MLC not present (not shown)
- Weak 0-1 km shear; stronger shear in 1-3 km layer
- Extreme lake-induced instability



(a)



(b)

As in previous figure except for a Type B event from 1200 UTC 8 January 2014.



Questions/Comments?



Joe.Villani@noaa.gov