


Meteorological Overview of the 28 January 2014 Southeast Winter Storm Trisha D. Palmer¹, S. Hunter Coleman², Patrick D. Moore³, Adam K. Baker¹, Jason T. Deese¹, Jessica L. Fieux¹, Steven E. Nelson¹, and Keith M. Stellman¹

Synoptic Overview

0000 UTC 28 January 2014 500 00 500 MB UA OBS, HGHTS, and TEMI 140128/1200 500 MB UA OBS, HGHTS, and TEMPS 850 Surfa

- Aloft
- Broad upper trough across eastern third of the country
- Upper divergence associated with the right entrance region of a jet streak
- Series of embedded shortwaves (90 knot wind maxima) contributed to deepening of mean upper trough
- Plains

¹National Weather Service Atlanta Forecast Office, Peachtree City, GA ²National Weather Service Forecast Office, Columbia, SC ³National Weather Service Greenville-Spartanburg Forecast Office, Greer, SC



years!

 Cold front moved across the Appalachian Mountains late on 27 January

• Temperatures dropped approximately 30°F in a 12hour period behind the front

• Temperatures continued to fall after onset of snow


SPC objective analysis of lower; dashed red, above

Climatological Reference Onset of snow with temperatures < 30°F *extremely* rare in the Southeast


• In Atlanta, it has only happened once in the past 30



- event, but road surfaces were initially still warm
- Widespread amounts of 1-2 inches of relatively dry snow Lower troposphere remained below freezing through the
- Initial snowfall melted on impact and subsequently re-froze • Frictional effects of tires on roads likely contributed to
- additional ice development Formation of thin layer of ice on streets and highways led to
- dangerous conditions
- Metro Atlanta was especially hard-hit; motorists stranded on roadways for hours and in some cases overnight







Accumulation and Impacts



Name: Trisha D. Palmer Organization: NOAA/NWS Peachtree City Email: trisha.palmer@noaa.gov