



NOAA

Overview

On the afternoon of April 13, 2014 a short lived tornado touched down within the community of Lovelady, TX injuring one and producing damage rated as EF-1 on the enhanced Fujita scale. This community lies in an area of poor low level radar coverage. At the reported time of the tornado, the radar presentation was of a weakening shower with no associated lightning. An analysis of upper air observations shows

that the tornado occurred in a region characterized by strong low level shear due to a strengthening low level jet and weak instability. Following this analysis an attempt is made to determine whether the tornado formed from a parent supercell or through non-supercell processes.

Radar Coverage



- Nearest WSR-88D to Lovelady is KHGX, 116 miles to the south
- Lowest elevation angle from KHGX (0.5°) is centered at 12,000ft over Lovelady
- Radar beamwidth is ~ 9,900ft (7,500ft to 17,500ft)



- 250mb analysis shows axis of upper level jet to the northwest of Lovelady and weakly diffluent flow across East Texas
- 500mb analysis shows a 50kt jet streak across East Texas (Lovelady falls into the right entrance region)
- •700mb analysis showing shortwave trough exiting the region (moved from NM to TX at approximate time of the tornado)
- 850mb analysis showing a strong low level jet extending southward into East Texas

Lovelady, TX: A Case Study of a Tornadic Cell in a Sparse Radar Coverage Environment Chris McKinney, Brian Haines, Scott Overpeck, and Dan Reilly National Weather Service Houston/Galveston, TX





- Est. time of tornado: 4:45 4:50 pm CDT (2145 UTC)
- First call to NWS Houston at 5:40 pm CDT
- Reflectivity weakening prior to tornado formation
- Little evidence of rotation from either radar, despite de-aliasing issues.
- Radar returns displaced south of Lovelady
- No watches or warnings for affected area

Synoptic Setup



- •Capping inversion present at 13/12z is removed 12 hours later
- Note strongly curved hodograph at 14/00z due to increasing low level winds
- •Strong low level jet present in KSHV sounding 55kts observed at 850mb level



- Surface analysis from 14/00z shows an approaching cold front across Eastern Oklahoma and North Texas
- Surface winds across East Texas are backed ahead of this feature
- High helicity values present across East Texas as a result of backed surface winds and strong low level jet









- 1 minor injury
- Trailer overturned and tree uprooted
- Damage rated as EF-1 on the Fujita Scale



- Requires pre-existing vertical vorticity
- Sharp wind shift along boundary
- Low CIN & high 0-3 km
- CAPE/ LCL
- layer shear
- thunderstorm life cycle

While the cause of the tornado cannot be definitively stated, evidence points toward development due to a low topped supercell: - Weak instability (CAPE 400 – 800 J/kg) in a highly sheared

- environment
- High SRH (SFC 1 km 400 500 m²s⁻²)
- Low LCL height
- free base, RFD clear slot

Detection of these events will remain problematic in areas with poor low level radar coverage.

References

Brady, R.H., and E.J. Szoke, 1989: A case study of nonmesocyclone tornado development in northeast Colorado: similarities to waterspout formation. Monthly Weather Review, 117, 843-856. Wakimoto, R. M., and J. W. Wilson, 1989: Non-supercell tornadoes. Mon. Wea. Rev., 117, 1113-1140.

Caruso, J. M., and J. M. Davies, 2005: Tornadoes in nonmesocyclone environments with pre-existing vertical vorticity along convergence boundaries. NWA Electronic J. Operational Meteorology, 2005-EJ4. [Available online at http://www.nwas.org/ej/.]

Burgess, D. W., R. J. Donaldson Jr., and P. R. Desrochers, 1993: Tornado detection and warning by radar. The Tornado: Its Structure, Dynamics, Prediction, and Hazards, Geophys. Monogr., No. 79, Amer. Geophys. Union, 203-221.

Wilczak, J.M., T.W. Christian, D.R. Wolfe, R.J. Zamora, and B. Stankov, Observations of a Colorado tornado, Part I, Mesoscale environment and tornadogenesis, Monthly Weather Review., 120, 497-520, 1992.

Acknowledgments

We thank Lance Wood for his helpful insights and suggestions.



Potential Tornado Formation Processes

Landspout

Steep low level lapse rates

• Small SRH and weak deep

Forms during updraft stage of

Low-topped Supercell



- Weak/moderate instability and low equilibrium level
- Strong shear ~40kts
- 0-3km storm relative helicity 200-500 m²s⁻²
- Bulk Richardson number 10 or less
- Forms during mature stage of thunderstorm

Conclusions

- No evidence of a pre-existing disturbance from earlier convection

- Visual appearance of tornado suggests parent supercell (rain

For more information contact: brian.haines@noaa.gov