

### III.7-MAP MEAN AREAL PRECIPITATION PROGRAM (MAP)

#### Bookmarks

Purpose . . . . .	<a href="#">[Bookmark]</a>
Program Input . . . . .	<a href="#">[Bookmark]</a>
Free Format Input Rules . . . . .	<a href="#">[Bookmark]</a>
Input Data . . . . .	<a href="#">[Bookmark]</a>
Program Execution Information . . . . .	<a href="#">[Bookmark]</a>
Sample Input and Output . . . . .	<a href="#">[Bookmark]</a>
Figures . . . . .	<a href="#">[Bookmark]</a>
<a href="#">[Bottom]</a>	

#### Contents

	<u>Page</u>
Purpose . . . . .	1
Program Input . . . . .	2
Free Format Input Rules . . . . .	2
Input Data . . . . .	3
Program Execution Information . . . . .	11
Sample Input and Output . . . . .	12

[\[Next\]](#) [\[Previous\]](#) [\[Bookmarks\]](#) [\[Top\]](#)

#### Purpose

The Mean Areal Precipitation (MAP) program provides techniques and procedures for computing estimates of mean areal precipitation from station precipitation data.

Chapter III.2 [\[Hyperlink\]](#) describes how to obtain the historical station data.

In most cases program Precipitation Preliminary Processing (PXPP) program (see Chapter III.7-PXPP [\[Hyperlink\]](#)) is used to analyze the station data before running program MAP.

Program MAP includes methods to:

1. Estimate missing precipitation amounts.
2. Distribute daily or accumulated precipitation into hourly amounts.
3. Adjust precipitation data for orographic and/or other effects by using mean monthly precipitation.
4. Create dummy (hypothetical) hourly or daily stations for which all precipitation data are estimated.

Program MAP includes the following options for calculating mean areal precipitation and generating output:

1. Compute grid point, Thiessen or input predetermined station weights. Grid point and Thiessen weights are computed using basin boundaries input as a series of latitude-longitude pairs.
2. Compute MAP in a 1, 3 or 6 hour data time interval.
3. Make consistency plots of a station's precipitation data without computing or printing MAP.
4. Generate consistency plots or specify predetermined station weights on a seasonal or on an annual basis. If the Interactive Double Mass Analysis (IDMA) program (see Section III.10-IDMA [[Hyperlink](#)]) is to be used to produce correction factors then the consistency plot option must be specified so that the input file needed by IDMA is created.
5. Print a summary of daily precipitation values that could not be distributed. The daily values are not distributed because all of the available hourly stations contain missing or zero precipitation for the 24 hour period. This situation occurs when localized events only occur where one or more daily stations are located. If large numbers of these cases occur then there may not be sufficient hourly stations in the analysis. If some daily stations have many more cases of values not being distributed than other daily stations then the observation times for those stations should be checked carefully.
6. Print the number of hours for which all hourly stations have missing data. Precipitation is set to zero for the hourly stations when this occurs. If this number is very large then additional hourly stations need to be added to the analysis.

Program MAP is currently dimensioned for the following maximum values:

- o 50 basins
- o 3000 basin boundary points per basin
- o 200 precipitation stations
- o 9 changes in observation time for each station
- o 10 precipitation adjustment factors for each station
- o 100 years of data
- o 3 groups of stations for consistency check

[\[Next\]](#) [\[Previous\]](#) [\[Bookmarks\]](#) [\[Top\]](#)

### Program Input

The program uses free format input for all cards except the Station Time Series Input File Information.

[\[Next\]](#) [\[Previous\]](#) [\[Bookmarks\]](#) [\[Top\]](#)

### Free Format Input Rules

See Chapter I.3-FREEFMT-UFIELD [[Hyperlink](#)] for a description of the general syntax rules.

Syntax rules specific to program MAP are:

1. The '@' indicates a card label. Card labels can be on the same line or the line above the fields of the card. For example:  
@A 10 1965 10 1966  
can be specified as  
@A  
10 1965 10 1966
2. If a card is not needed for a particular run then the label and all fields must be omitted.
3. Not all letters are used as card labels. For example there is no card K.
4. Character fields containing commas and embedded blanks must be enclosed by apostrophes. Note that filenames (card A field 9) and area descriptions (card I field 2) generally contain blanks.
5. All fields are required. A null field (double commas) must be used to denote single fields for which default values are to be used. If N consecutive fields use default values then N+1 commas must be used. The following input implies that defaults are to be used for the first two fields on card B:  
@B , , ,
6. Not all fields have valid defaults. If the documentation does not specify a default then the input must be specified.
7. Consecutive commas at the end of a card can be omitted. For example:  
@C GRID , , ,  
can be specified as  
@C GRID

[\[Next\]](#) [\[Previous\]](#) [\[Bookmarks\]](#) [\[Top\]](#)

### Input Data

Card   Field   Format   Contents

#### **PERIOD OF RECORD AND OUTPUT OPTIONS**

A	1		'@A'
	2	I	First month
	3	I	First year (four digits)
	4	I	Last month
	5	I	Last year (four digits)
	6	A	Units in which MAP is to be calculated and printed; allowable units are inches ('IN') or

<u>Card</u>	<u>Field</u>	<u>Format</u>	<u>Contents</u>
			millimeters ('MM'); default is 'IN'
7	A		Units in which MAP results will be written into output file ('IN' or 'MM'); default is 'IN'
8			Not used; enter a null field
9	A		Option to list MAP results; default is to not list (MAP results must be written to a file in order to be listed): 'LIST' = list MAP time series results
10	A20		Forecast Group directory name; this will become part of the directory name into which MAP results will be written; embedded blanks are not allowed <u>1/</u> [ <u>Bookmark#1</u> ] <u>2/</u> [ <u>Bookmark#1</u> ]  Leave this field blank if MAP results are not to be written to a file

#### **WEIGHTING OPTIONS**

B	1		'@B'
	2	I	Number of areas (enter zero to have only consistency plots made; MAP will not be computed or printed)
	3	A	Station weighting options; default is 'THIE': 'GRID' = compute grid point station weights for each area 'THIE' = compute Thiessen station weights for each area 'PRE' = read predetermined annual station weights 'SEAS' = read predetermined seasonal station weights for each area (must specify first month of winter in field 8 and the first month of summer in field 9 on this card)
	4		Not used; enter a null field
	5	A	Stop option; default is 'CONT': 'STOP' = terminate program after printing station weights 'CONT' = continue program to calculate MAP
	6	I	MAP time series data time interval (1, 3 or 6 hours); default is 6 hours

<u>Card</u>	<u>Field</u>	<u>Format</u>	<u>Contents</u>
	7	A	Precipitation adjustment option; default is no precipitation adjustment; adjustments are entered on card O: 'ADJ' = precipitation adjustments required
	8	A	Consistency plot option; default is no consistency plot; consistency plot information is entered on cards R and S; this option must be specified if the IDMA program is to be used to determine correction factors: 'PLOT' = develop precipitation consistency plots 'SESN' = develop precipitation consistency plots on a seasonal basis

If 'SESN' is specified as consistency plot option or 'SEAS' is specified as station weighting option then enter beginning winter and summer months:

	9	I	First month of winter; default is 10
	10	I	First month of summer; default is 5
	11	A	Observation time correction option; default is no time corrections; corrections are entered on card M: 'CTIM' = time corrections to be read

#### **INPUT AND OUTPUT OPTIONS**

C	1		'@C'
	2	A	Mean monthly precipitation option; default is to not use station means: 'NORM' = use mean (normal) monthly precipitation when estimating missing data
	3	A	MAP output option; default is to print MAP summary and table of distribution messages: 'COMP' = print all distribution messages (contact your Focal Point first because this creates large amounts of output)
	4	A	Precipitation comparison option; default is no comparison: 'CMPR' = precipitation comparison desired for evaluation of estimation procedure
	5	R	Minimum station weight; default is 0.01

Card   Field   Format   Contents

**STATION INFORMATION**

D	1		'@D'
	2	I	Number of hourly stations <u>3/</u> [ <u>Bookmark#1</u> ]
	3	I	Number of daily stations <u>3/</u> [ <u>Bookmark#2</u> ]

Repeat cards F and G for each precipitation station.

The order in which the stations are read determines the run sequence numbers.

F	1		'@F'
	2	A20	Station name
	3	R,I, I-I	Station latitude <u>4/</u> [ <u>Bookmark#1</u> ]
	4	R,I, I-I	Station longitude <u>4/</u> [ <u>Bookmark#2</u> ]
	5	R	Observation time for initial month in local standard time (hours 1 through 24) 0 = hourly station
	6	A7	Station identifier or 'DUMMY' if dummy station; default is non-dummy station <u>14/</u> [ <u>Bookmark</u> ]

G	1		'@G'
	2-13	R	Mean monthly precipitation values (only used if 'NORM' is specified on card C)

**PRECIPITATION COMPARISON**

Card H is needed only if precipitation comparison are to be done (field 3 on card C).

H	1		'@H'
	2	I	Sequence number of station to be replaced
	3	I	Sequence number of dummy station

**AREA INFORMATION AND PREDETERMINED WEIGHTS**

Omit cards I, J and L if number of areas on card B is zero.

Repeat cards I, J and L for each area in the run.

<u>Card</u>	<u>Field</u>	<u>Format</u>	<u>Contents</u>
I	1		'@I'
	2	A12	Area identifier
	3	A20	Area description
	4	R	Area size in square kilometers or square miles
	5	A	Area units: 'KM2' = square kilometers 'MI2' = square miles
	6	A12	Basin name (optional); if specified then this name will become part of the directory name and file name into which MAP results will be written (./basin_name/basin_name.MAPnn); if omitted then the area identifier is used as part of the filename and the additional directory level is omitted (./area_id.MAPnn) <u>1/ [Bookmark#2]</u> <u>2/ [Bookmark#2]</u>
	7	A12	File name (optional); if specified then the MAP output will be written to that file (./basin_name/file_name.MAPnn); if omitted then the file name will be set to the basin name <u>1/ [Bookmark#3]</u> <u>2/ [Bookmark#3]</u>

Omit card J if using predetermined weights.

The format of card J is the same as for the PPINIT program DEFINE BASIN command. 5/ [Bookmark]

J	1		'@J'
	2	A	'BASE' or 'BASE(units)'; units can be 'METR' for metric or 'ENGL' for English; default is 'ENGL'
	3	A8	Basin identifier
	4	A20	Descriptive information
	5	(R,R)	Latitude/longitude pairs defining basin boundary <u>4/ [Bookmark#3]</u> <u>6/ [Bookmark]</u> <u>7/ [Bookmark]</u> or Latitude/longitude pair defining basin centroid <u>8/ [Bookmark]</u>

Omit card L if not using predetermined weights.

L	1		'@L'
---	---	--	------

<u>Card</u>	<u>Field</u>	<u>Format</u>	<u>Contents</u>
	2-n	R	Predetermined station weights (order determined from card F)
			If not using seasonal weights then enter annual station weight for each station
			If using seasonal weights then enter winter station weights for all stations followed by summer weights for all stations)

### **OBSERVATION TIME CHANGES**

Card M is needed only if changes are to be made to the observation times (field 10 on card B).

Changes in observation time for non-recording stations. One change per card. Maximum of 9 changes per station.

Repeat card M for each change.

M	1		'@M'
	2	I	Station sequence number Enter '999' for end of card group <u>9</u> / [ <u>Bookmark#1</u> ]
	3	I	Month of time change
	4	I	Year (four digits)
	5	R	New observation time in hours (1 to 24)

### **PRECIPITATION ADJUSTMENTS**

Card O is needed only if adjustments are to be made to the precipitation data (field 6 on card B). 10/ [Bookmark]

One change per card with a maximum of 10 changes per station.

Repeat card O for each precipitation adjustment factor.

O	1		'@O'
	2	A	Season for which precipitation adjustment factor is to be applied; default is winter and summer: 'W' = winter 'S' = summer Enter '999' for end of card group <u>9</u> / [ <u>Bookmark#2</u> ]
	3	I	Sequence number of station





The full pathname is created by concatenating the Forecast Group directory name to a pathname generated by .Apps\_defaults tokens (see Chapter I.2-UNIX-CALB [[Hyperlink](#)]). The directory defined by this full pathname must be created by the user.

The Forecast Group directory name can be specified as:

name1  
or  
name1/name2

- [\[Back#1\]](#) [\[Back#2\]](#) [\[Back#3\]](#)
- 2/ The characters 'MAPnn' are appended to the basin name or area identifier to create the full file name where nn is the data time interval. The maximum length of the full pathname plus the file name is 112 characters.
- [\[Back#1\]](#) [\[Back#2\]](#) [\[Back#3\]](#)
- 3/ This number must include all dummy stations.
- [\[Back#1\]](#) [\[Back#2\]](#)
- 4/ The station location can be input in decimal degrees or degrees and minutes. For example specifying 50.5 indicates decimal degrees and is equivalent to 50-30 for 5030 which indicates degrees and minutes. Valid values for minutes are 0 to 59.
- If degrees and minutes are input and the optional '-' is omitted then two characters must always be used to specify the minutes. For example 5003 would be a valid input field to specify 50 degrees and 3 minutes.
- [\[Back#1\]](#) [\[Back#2\]](#) [\[Back#3\]](#)
- 5/ See Section VI.3.3B-DEFINE-BASIN [[Hyperlink](#)] for a description of the input data. The first three and last two optional fields are not used by program MAP. These fields do not need to be included. If they are included then they will be ignored.
- [\[Back\]](#)
- 6/ Pairs must be entered in clockwise order and in decimal degrees. The first value must be preceded by an opening parenthesis. The last value must be followed by a closing parenthesis.
- [\[Back\]](#)
- 7/ The basin boundary must have at least 3 pairs.
- [\[Back\]](#)
- 8/ If only one latitude/longitude pair is entered it is assumed to be the basin centroid. The pair must be enclosed in parentheses.
- [\[Back\]](#)
- 9/ The '999' signifies the end of the data. The field following the '999' must contain a card label.
- [\[Back#1\]](#) [\[Back#2\]](#)
- 10/ Adjustments to each station must be in order by date (earliest first). Adjustments remain in effect until the date of the next correction.
- [\[Back\]](#)
- 11/ Date must be within the observed period for the station. If it is not then the correction will be ignored.
- [\[Back#1\]](#) [\[Back#2\]](#)
- 12/ The Q card must be on a separate line preceding the Station Time Series Input File Information cards.

One filename must be included for each station to be used. The maximum number of characters in the filename is 40. See Chapter I.2-UNIX-CALB [[Hyperlink](#)] for a description of the directory structure.

No Time Series Input File Information card is needed for a DUMMY station.

The Time Series File Input Information cards for hourly stations must be entered first followed by the Time Series File Input Information cards for daily stations. The Time Series File Input Information cards are entered in the same station order used for entering cards F and G.

The period of record for each time series can be different and need not coincide with the MAP period of record to be calculated. Time series data are read according to the following rules:

- o If the time series begins before or ends after the desired MAP period of record then the extra months are ignored.
- o If the time series begins after or ends before the desired MAP period of record then missing data symbols are written for the missing months.

[[Back](#)]

13/ Stations with positive sequence numbers make up the group base and each is plotted against the average of the other base stations. Base must contain at least two stations. Each station with a negative sequence number is dropped from the average of the group base stations and is plotted against the group base average. Station order does not matter.

For example group one contains 5 stations. Sequence numbers are 3, 5, 6, 9 and 12. Card S input for group one is:

```
6 5 9 -12 -3
```

The group base consists of stations 6, 5 and 9. Station 6 is plotted against the average of stations 5 and 9, station 5 against the average of 6 and 9, station 9 against the average of 5 and 6. Stations 12 and 3 are each plotted against the average of the group base stations.

[[Back](#)]

14/ The station identifier is used by program IDMA and is entered as the state and station number. The state can be entered as the 2 character state abbreviation or the NCDC state number (see Chapter I.6 [[Hyperlink](#)]). For example AL0001 and 010001 would be entered for station 0001 for Alabama.

[[Back](#)]

[[Next](#)] [[Previous](#)] [[Bookmarks](#)] [[Top](#)]

### Program Execution Information

See Chapter I.2 [[Hyperlink](#)] for program execution information.

[\[Next\]](#) [\[Previous\]](#) [\[Bookmarks\]](#) [\[Top\]](#)

Sample Input and Output

Figure 1 [\[Bookmark\]](#) is sample program input.

Figure 2 [\[Bookmark\]](#) is sample program output.

Figure 1. Sample input for program MAP

```
$ SAMPLE INPUT FOR PROGRAM MAP

@A 10 1959 9 1961 IN IN ,, LIST map_ts
@B 2 GRID 1 CONT 6 ADJ PLOT
@C ''
@D 5 5

@F 'QUEBEC, NC' 35.17 82.89 0.0
@F 'GLOUCESTER GAP,NC' 35.27 82.84 0.0
@F 'PINK BEDS,NC' 35.33 82.78 0.0
@F 'CEDAR MTN,NC' 35.15 83.65 0.0
@F 'HAYWOOD GAP,NC' 35.30 82.91 0.0
@F 'BREVARD, NC' 35.23 82.73 18.
@F 'OWENS GAP,NC' 35.20 82.98 7.
@F 'PISGAH FOREST 1N,NC' 35.27 82.70 8.
@F 'ROSMAN, NC' 35.24 82.83 8.
@F 'SASSAFRASS MTS, SC' 35.06 82.77 8.

@I MAP-0343990 'FRENCH BROAD, ROSMAN' 67.9 MI2 french_broad
@J (35.13,82.82 35.13,82.84 35.12,82.84 35.13,82.88 35.10,82.90
 35.17,82.93 35.16,82.97 35.19,82.98 35.20,82.92 35.22,82.92
 35.23,82.93 35.30,82.92 35.31,82.88 35.27,82.87 35.26,82.85)
@I MAP-0344300 'BLANTYRE LOCAL' 188. MI2 blantyre
@J (35.30,82.63 35.26,82.61 35.26,82.56 35.20,82.54 35.16,82.59
 35.14,82.58 35.12,82.67 35.13,82.69 35.10,82.70 35.06,82.77
 35.10,82.81 35.00,82.83 35.12,82.84 35.13,82.84 35.13,82.82
 35.26,82.85 35.26,82.80 35.28,82.74 35.27,82.70 35.33,82.74)

@O ,, 9 7 1960 .87
@O 999

@Q
Quebec_NC
GloucesterGap_NC
PinkBeds_NC
CedarMtn_NC
HaywoodGap_NC
Brevard_NC
OwensGap_NC
PisgahForest1N_NC
Rosman_NC
SassafrassMtn_SC

@R 1 10

@S 1 2 3 4 5 6 7 8 9 10
```





Figure 2. Sample output from program MAP

28	.00	.00	.00	.00 /	.00	.21	.44	.19 /	.22	.25	.17	.22 /
31	.04	.06	.00	.00 /								
11/1959												
1	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
4	.00	.00	.00	.00 /	.04	.02	.08	.31 /	.22	.00	.00	.00 /
7	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
10	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
13	.00	.00	.00	.00 /	.09	.05	.00	.00 /	.05	.05	.00	.00 /
16	.00	.00	.05	.23 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
19	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.01	.02	.00	.00 /
22	.00	.00	.00	.00 /	.00	.00	.02	.03 /	.96	.17	.01	.00 /
25	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.69 /
28	.01	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.01 /
12/1959												
1	.00	.00	.00	.00 /	.00	.00	.14	.04 /	.02	.01	.00	.00 /
4	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.11	.27	.03	.02 /
7	.01	.01	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
10	.00	.00	.00	.00 /	.00	.00	.01	.23 /	.73	.76	.06	.06 /
13	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
16	.00	.00	.00	.00 /	.00	.03	.04	.15 /	.22	.64	.30	.10 /
19	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
22	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.05 /
25	.12	.03	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.02	.11 /
28	.52	.83	.13	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
31	.00	.00	.00	.00 /								
[Time series data listing for months 1/1960 through 6/1961 deleted.]												
7/1961												
1	.00	.00	.03	.00 /	.00	.00	.12	.00 /	.00	.00	.00	.00 /
4	.01	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
7	.00	.22	.31	.02 /	.00	.00	.00	.00 /	.00	.02	.00	.00 /
10	.00	.00	.00	.00 /	.00	.00	.00	.44 /	.25	.07	.25	.08 /
13	.01	.02	.33	.00 /	.00	.23	.00	.01 /	.00	.01	.29	.00 /
16	.00	.01	.28	.11 /	.01	.09	.00	.00 /	.00	.01	.21	.01 /
19	.00	.04	.05	.03 /	.00	.00	.03	.42 /	.01	.03	.19	.00 /
22	.00	.17	.18	.03 /	.00	.19	.31	.00 /	.00	.00	.00	.03 /
25	.00	.00	.00	.00 /	.00	.00	1.75	.02 /	.00	.00	.09	.00 /
28	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.08	.08	.00 /
31	.00	.10	.00	.00 /								
8/1961												
1	.00	.01	.00	.00 /	.00	.01	.15	.15 /	.00	.03	.23	.00 /
4	.00	.03	.10	.93 /	.01	.02	.00	.00 /	.00	.00	.04	.00 /
7	.00	.02	1.09	.36 /	.00	.05	.10	.00 /	.00	.01	.52	1.91 /
10	.47	.00	.01	.00 /	.00	.01	.00	.00 /	.00	.02	.00	.00 /
13	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
16	.00	.00	.00	.00 /	.00	.03	.00	.00 /	.00	.00	.01	.33 /
19	.01	.00	.00	.00 /	.00	.00	.01	.00 /	.00	.07	1.05	.34 /
22	.00	.03	.20	.06 /	.00	.04	.00	.48 /	2.60	.66	1.96	.07 /
25	.43	.93	.20	.00 /	.00	.08	.21	.10 /	.03	.06	.01	.86 /
28	.16	.07	.01	.00 /	.00	.00	.06	.18 /	.00	.05	.19	.06 /
31	.00	.02	.00	.04 /								
9/1961												
1	.01	.02	.37	.00 /	.00	.00	.00	.00 /	.00	.01	.02	.00 /
4	.00	.37	.00	.00 /	.00	.00	.04	.00 /	.00	.00	.00	.00 /
7	.17	.05	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
10	.00	.00	.00	.01 /	.35	.08	.10	.00 /	.00	.00	.00	.00 /
13	.00	.01	.00	.00 /	.00	.05	.02	.02 /	.00	.00	.00	.00 /
16	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.15	.02 /
19	.11	.20	.09	.03 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
22	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
25	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
28	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
- LISTING OF DATA (UNITS=IN )FOR THE TIME SERIES BLANTYRE LOCAL												
10/1959												
1	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
4	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.01	.53 /
7	.44	.08	.03	.01 /	.25	.39	.48	.62 /	1.06	.56	.01	.13 /
10	.11	.21	.50	.17 /	.07	.01	.00	.00 /	.00	.00	.00	.00 /
13	.00	.00	.02	.52 /	1.25	.28	.15	.00 /	.00	.01	.00	.02 /
16	.00	.00	.00	.00 /	.01	.00	.34	.03 /	.01	.01	.00	.00 /
19	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.01	.03	.11 /
22	.13	.02	.00	.00 /	.07	.24	.64	.05 /	.05	.04	.00	.00 /
25	.00	.01	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
28	.00	.00	.00	.00 /	.00	.23	.24	.10 /	.14	.27	.17	.26 /
31	.05	.07	.00	.00 /								
11/1959												
1	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
4	.00	.00	.00	.00 /	.03	.03	.04	.10 /	.07	.00	.00	.00 /
7	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
10	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /
13	.00	.00	.00	.00 /	.02	.03	.00	.00 /	.04	.07	.00	.00 /
16	.00	.00	.02	.14 /	.00	.00	.00	.00 /	.00	.00	.01	.00 /
19	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.01	.00	.00 /
22	.00	.00	.00	.00 /	.00	.01	.05	.04 /	1.21	.15	.01	.00 /
25	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.46 /
28	.00	.00	.00	.00 /	.00	.00	.00	.00 /	.00	.00	.00	.00 /





Figure 2. Sample output from program MAP

12/1960	85.2	92.6	98.7	91.0	88.2	92.2	96.1	91.3	101.1	90.8
1/1961	87.3	95.8	102.4	94.2	90.8	95.5	100.3	94.4	105.2	93.9
2/1961	98.2	106.5	113.2	104.9	101.2	106.2	112.7	104.9	119.2	104.2
3/1961	104.4	113.8	121.2	111.9	108.4	113.3	119.2	112.1	128.7	111.0
4/1961	110.1	119.4	126.7	117.6	113.2	119.1	125.8	117.7	135.1	116.6
5/1961	113.7	124.7	131.9	122.7	118.1	124.3	132.0	122.7	141.9	121.6
6/1961	122.5	134.7	142.1	132.6	125.5	134.4	144.5	132.3	151.2	131.6
7/1961	130.0	141.9	147.8	139.9	131.7	141.7	151.6	139.5	158.2	138.8
8/1961	149.2	159.5	166.0	157.6	148.6	159.6	172.0	157.0	172.6	156.9
9/1961	151.7	161.6	167.7	159.8	150.8	161.7	174.2	159.1	174.8	159.1

DEVIATION OF ACCUMULATED PRECIPITATION FROM THE GROUP BASE-----GROUP=1

STA. RUN NO.	STA. PLOT NO.	STATION NAME
1	1	QUEBEC, NC
2	2	GLOUCESTER GAP, NC
3	3	PINK BEDS, NC
4	4	CEDAR MTN, NC
5	5	HAYWOOD GAP, NC

ASTERISKS INDICATE ZERO DEVIATION

UNITS ARE IN

BASE PX	-39.6	-33.6	-27.6	-21.6	-15.6	-9.6	-3.6	2.4	8.4	14.4	20.4
.0	.	.	.	.	.	.	*	.	.	.	.
1.7	.	.	.	.	.	.	*	.	.	.	.
3.4	.	.	.	.	.	.	*	.	.	.	.
5.1	.	.	.	.	.	.	*	.	.	.	.
6.8	.	.	.	.	.	.	*	.	.	.	.
8.5	.	.	.	.	.	.	*	.	.	.	.
10.2	.	.	.	.	.	.	*	.	.	.	.
11.9	.	.	.	.	.	.	*	.	.	.	.
13.6	.	.	.	.	.	.	1 *	35.4	.	.	.
15.3	.	.	.	.	.	.	*	245	.	.	.
17.0	.	.	.	.	.	.	1 *	3	.	.	.
18.7	.	.	.	.	.	.	*	.	.	.	.
20.4	.	.	.	.	.	.	*	.	.	.	.
22.1	.	.	.	.	.	.	1 *	3 2 45	.	.	.
23.8	.	.	.	.	.	.	*	.	.	.	.
25.5	.	.	.	.	.	.	*	.	.	.	.
27.2	.	.	.	.	.	.	*	.	.	.	.
28.9	.	.	.	.	.	.	1 *	3 4 5	.	.	.
30.6	.	.	.	.	.	.	*	.	.	.	.
32.3	.	.	.	.	.	.	*	.	.	.	.
34.0	.	.	.	.	.	.	*	.	.	.	.
35.7	.	.	.	.	.	.	*	.	.	.	.
37.4	.	.	.	.	.	.	3	2 4 5	.	.	.
39.1	.	.	.	.	.	.	1 *	.	.	.	.
40.8	.	.	.	.	.	.	*	.	.	.	.
42.5	.	.	.	.	.	.	*	.	.	.	.
44.2	.	.	.	.	.	.	*	.	.	.	.
45.9	.	.	.	.	.	.	3	2 54	.	.	.
47.6	.	.	.	.	.	.	1 *	.	.	.	.
49.3	.	.	.	.	.	.	*	5 4	.	.	.
51.0	.	.	.	.	.	.	1 3*	2	.	.	.
52.7	.	.	.	.	.	.	*	4	.	.	.
54.4	.	.	.	.	.	.	1 3*	2 5	.	.	.
56.1	.	.	.	.	.	.	*	.	.	.	.
57.8	.	.	.	.	.	.	*	.	.	.	.
59.5	.	.	.	.	.	.	*	5 2	.	.	.
61.2	.	.	.	.	.	.	1 3	4	.	.	.
62.9	.	.	.	.	.	.	*	.	.	.	.
64.6	.	.	.	.	.	.	3 *	4 2 5	.	.	.
66.3	.	.	.	.	.	.	1 *	.	.	.	.
68.0	.	.	.	.	.	.	*	.	.	.	.
69.7	.	.	.	.	.	.	*	.	.	.	.
71.4	.	.	.	.	.	.	*	.	.	.	.
73.1	.	.	.	.	.	.	*	.	.	.	.
74.8	.	.	.	.	.	.	3 *	425	.	.	.
76.5	.	.	.	.	.	.	1 *	.	.	.	.
78.2	.	.	.	.	.	.	*	.	.	.	.
79.9	.	.	.	.	.	.	*	4 25	.	.	.
81.6	.	.	.	.	.	.	1 3	*	.	.	.
83.3	.	.	.	.	.	.	*	.	.	.	.
85.0	.	.	.	.	.	.	*	.	.	.	.
86.7	.	.	.	.	.	.	3 *	4 2 5 5	.	.	.
88.4	.	.	.	.	.	.	11 3	4 2	.	.	.
90.1	.	.	.	.	.	.	*	.	5	.	.
91.8	.	.	.	.	.	.	1 3	4 2	5	.	.
93.5	.	.	.	.	.	.	*	.	.	.	.
95.2	.	.	.	.	.	.	1 3	4	.	.	.
96.9	.	.	.	.	.	.	*	.	.	.	.
98.6	.	.	.	.	.	.	*	.	.	.	.
100.3	.	.	.	.	.	.	*	.	.	.	.
102.0	.	.	.	.	.	.	*	.	.	.	.
103.7	.	.	.	.	.	.	*	.	.	.5	.
105.4	.	.	.	.	.	.	3 *	42	.	.	.
107.1	.	.	.	.	.	.	1 *	.	.	.	.
108.8	.	.	.	.	.	.	*	.	.	.	.
110.5	.	.	.	.	.	.	*	.	.	5	.
112.2	.	.	.	.	.	.	*	4 2	.	.	.
113.9	.	.	.	.	.	.	1 3	*	.	.	.
115.6	.	.	.	.	.	.	*	.	.	.	.
117.3	.	.	.	.	.	.	*	42	.	5	.
119.0	.	.	.	.	.	.	1 3	*	.	.	.

Figure 2. Sample output from program MAP

120.7	.	.	.	.	.	.	.	*	.	.	.	.
122.4	.	.	.	.	.	.	.	*	.	.	.	.
124.1	.	.	.	.	.	1	3	*	.	.	.	.
125.8	.	.	.	.	.	.	.	*	.	.	.	.
127.5	.	.	.	.	.	.	.	*	.	.	.	.
129.2	.	.	.	.	.	.	.	*	.	.	.	.
130.9	.	.	.	.	.	.	.	*	.	.	.	5.
132.6	.	.	.	.	.	.	.	*	.	.	.	.
134.3	.	.	.	.	.	1	.3	*	.	.	.	.
136.0	.	.	.	.	.	.	.	*	.	.	.	.
137.7	.	.	.	.	.	.	.	*	.	.	.	.
139.4	.	.	.	.	.	.	.	*	.	.	.	.
141.1	.	.	.	.	.	1	3.	*	.	2.	4	5
142.8	.	.	.	.	.	.	.	*	.	.	.	.
144.5	.	.	.	.	.	.	.	*	.	.	.	.
146.2	.	.	.	.	.	.	.	*	.	.	.	.
147.9	.	.	.	.	.	.	.	*	.	.	.	.
149.6	.	.	.	.	.	.	.	*	.	.	.	.
151.3	.	.	.	.	.	.	.	*	.	.	.	.
153.0	.	.	.	.	.	.	.	*	.	.	.	.
154.7	.	.	.	.	.	.	.	*	.	.	.	.
156.4	.	.	.	.	.	.	.	*	.	.	.	.45
158.1	.	.	.	.	.	.	.	*	.	2	.	.
159.8	.	.	.	.	.	31.	.	*	.	2.	.45	.
161.5	.	.	.	.	.	3	1	*	.	.	.	.
163.2	.	.	.	.	.	.	.	*	.	.	.	.
164.9	.	.	.	.	.	.	.	*	.	.	.	.
166.6	.	.	.	.	.	.	.	*	.	.	.	.
168.3	.	.	.	.	.	.	.	*	.	.	.	.
170.0	.	.	.	.	.	.	.	*	.	.	.	.

MONTHLY LISTING OF ACCUMULATED PRECIPITATION AND BASE PRECIPITATION.

MO/YEAR	BREVARD, NC		OWENS GAP, NC		PISGAH FOREST 1N, NC		ROSMAN, NC		SASSAFRASS MTS, SC	
	ACC. PX	BASE PX	ACC. PX	BASE PX	ACC. PX	BASE PX	ACC. PX	BASE PX	ACC. PX	BASE PX
10/1959	9.8	13.6	14.8	13.0	9.2	13.7	10.7	13.5	16.2	12.9
11/1959	12.0	16.8	17.6	16.2	11.4	16.8	14.5	16.5	19.2	16.0
12/1959	16.8	22.6	22.5	21.9	15.9	22.7	22.7	21.9	24.2	21.8
1/1960	22.8	29.6	28.6	29.0	22.3	29.7	29.8	28.9	30.3	28.8
2/1960	31.3	39.0	37.5	38.3	31.5	39.0	39.9	38.0	40.1	38.0
3/1960	38.3	47.2	47.7	46.2	39.5	47.1	47.9	46.2	48.1	46.1
4/1960	42.2	51.4	50.6	50.5	43.2	51.3	52.5	50.2	53.0	50.2
5/1960	45.7	55.2	54.0	54.3	46.5	55.1	57.3	53.9	57.2	53.9
6/1960	53.3	61.8	59.1	61.2	53.5	61.8	62.1	60.8	63.9	60.6
7/1960	57.6	66.1	63.0	65.5	58.1	66.0	65.9	65.2	71.4	64.6
8/1960	65.7	76.1	71.2	75.5	64.3	76.3	78.4	74.7	84.5	74.0
9/1960	70.4	81.8	78.4	80.9	67.9	82.1	85.1	80.1	91.2	79.5
10/1960	77.9	88.2	85.3	87.4	75.6	88.5	90.5	86.8	97.8	86.0
11/1960	78.7	89.5	86.5	88.6	76.3	89.8	91.7	88.0	99.6	87.2
12/1960	82.0	92.9	89.1	92.1	79.5	93.2	95.1	91.4	103.1	90.6
1/1961	84.6	96.1	92.1	95.3	82.3	96.4	97.9	94.7	107.0	93.7
2/1961	93.9	107.0	102.9	106.0	91.2	107.3	107.9	105.5	116.7	104.5
3/1961	100.0	114.2	110.8	113.0	97.0	114.6	113.9	112.7	124.6	111.5
4/1961	105.6	119.9	115.9	118.8	102.0	120.3	119.7	118.3	130.7	117.1
5/1961	110.0	125.2	119.4	124.1	107.0	125.5	127.0	123.3	135.5	122.3
6/1961	119.5	135.1	129.0	134.0	115.0	135.6	138.8	132.9	147.1	132.0
7/1961	128.1	142.1	133.1	141.6	119.8	143.0	148.9	139.8	157.9	138.8
8/1961	149.0	159.5	144.3	160.0	138.5	160.7	173.8	156.8	170.7	157.1
9/1961	151.3	161.7	146.5	162.2	140.2	162.9	175.9	158.9	173.4	159.2

DEVIATION OF ACCUMULATED PRECIPITATION FROM THE GROUP BASE----GROUP=1

STA. RUN NO.	STA. PLOT NO.	STATION NAME
6	1	BREVARD, NC
7	2	OWENS GAP, NC
8	3	PISGAH FOREST 1N, NC
9	4	ROSMAN, NC
10	5	SASSAFRASS MTS, SC

ASTERISKS INDICATE ZERO DEVIATION UNITS ARE IN

BASE PX	-40.8	-34.8	-28.8	-22.8	-16.8	-10.8	-4.8	1.2	7.2	13.2	19.2
.0	.	.	.	.	.	.	.	*	.	.	.
1.7	.	.	.	.	.	.	.	*	.	.	.
3.4	.	.	.	.	.	.	.	*	.	.	.
5.1	.	.	.	.	.	.	.	*	.	.	.
6.8	.	.	.	.	.	.	.	*	.	.	.
8.5	.	.	.	.	.	.	.	*	.	.	.
10.2	.	.	.	.	.	.	.	*	.	.	.
11.9	.	.	.	.	.	.	.	*	.	.	.
13.6	.	.	.	.	.	.	.	*	.	.	.
15.3	.	.	.	.	.	.	.	*	.	.	.
17.0	.	.	.	.	.	.	.	*	.	.	.
18.7	.	.	.	.	.	.	.	*	.	.	.
20.4	.	.	.	.	.	.	.	*	.	.	.
22.1	.	.	.	.	.	.	.	*	.	.	.
23.8	.	.	.	.	.	.	.	*	.	.	.
25.5	.	.	.	.	.	.	.	*	.	.	.
27.2	.	.	.	.	.	.	.	*	.	.	.
28.9	.	.	.	.	.	.	.	*	.	.	.
30.6	.	.	.	.	.	.	.	*	.	.	.
32.3	.	.	.	.	.	.	.	*	.	.	.
34.0	.	.	.	.	.	.	.	*	.	.	.

Figure 2. Sample output from program MAP

35.7	.	.	.	.	.	.	.	.	.	*	.	.	.
37.4	.	.	.	.	.	.	.	.	.	*	.5	.	.
39.1	.	.	.	.	.	.	3	.	.	2*	.	.	.
40.8	.	.	.	.	.	.	.	.	.	*	.	.	.
42.5	.	.	.	.	.	.	.	.	.	*	.	.	.
44.2	.	.	.	.	.	.	.	.	.	*	.	.	.
45.9	.	.	.	.	.	.	.	.	.	*	.5	.	.
47.6	.	.	.	.	.	.	1	3	.	*	.	.	.
49.3	.	.	.	.	.	.	.	.	.	*	.	.	.
51.0	.	.	.	.	.	.	13	.	.	2	.45	.	.
52.7	.	.	.	.	.	.	.	.	.	*	.	.	.
54.4	.	.	.	.	.	.	1	3	.	2*	.54	.	.
56.1	.	.	.	.	.	.	.	.	.	*	.	.	.
57.8	.	.	.	.	.	.	.	.	.	*	.	.	.
59.5	.	.	.	.	.	.	.	.	.	*	.	.	.
61.2	.	.	.	.	.	.	3	.	2	*4	5	.	.
62.9	.	.	.	.	.	.	.	.	.	*	.	.	.
64.6	.	.	.	.	.	.	.	.	.	*	.	.	.
66.3	.	.	.	.	.	.	13	.	2	*4	5	.	.
68.0	.	.	.	.	.	.	.	.	.	*	.	.	.
69.7	.	.	.	.	.	.	.	.	.	*	.	.	.
71.4	.	.	.	.	.	.	.	.	.	*	.	.	.
73.1	.	.	.	.	.	.	.	.	.	*	.	.	.
74.8	.	.	.	.	.	.	.	.	.	*	.	.	.
76.5	.	.	.	.	.	.	3	.1	.	*	4	5	.
78.2	.	.	.	.	.	.	.	.	.	*	.	.	.
79.9	.	.	.	.	.	.	.	.	.	*	4	5	.
81.6	.	.	.	.	.	3	1	.	2	*	.	.	.
83.3	.	.	.	.	.	.	.	.	.	*	.	.	.
85.0	.	.	.	.	.	.	.	.	.	*	.	.	.
86.7	.	.	.	.	.	.	.	.	2	*	4	55	.
88.4	.	.	.	.	.	3	1	.	2	*	4	.	.
90.1	.	.	.	.	.	3	1	.	.	*	.	5	.
91.8	.	.	.	.	.	.	.	.	2	*	4	.	.
93.5	.	.	.	.	.	3	1	.	.	*	.	5	.
95.2	.	.	.	.	.	.	.	.	2	*	4	.	.
96.9	.	.	.	.	.	3	1	.	.	*	.	.	.
98.6	.	.	.	.	.	.	.	.	.	*	.	.	.
100.3	.	.	.	.	.	.	.	.	.	*	.	.	.
102.0	.	.	.	.	.	.	.	.	.	*	.	.	.
103.7	.	.	.	.	.	.	.	.	.	*	.	5	.
105.4	.	.	.	.	.	.	.	.	2	*	4	.	.
107.1	.	.	.	.	.	.3	1	.	.	*	.	.	.
108.8	.	.	.	.	.	.	.	.	.	*	.	.	.
110.5	.	.	.	.	.	.	.	.	.	*	.	.	.
112.2	.	.	.	.	.	.	.	.	2	*	4	5	.
113.9	.	.	.	.	.	3	1	.	.	*	.	.	.
115.6	.	.	.	.	.	.	.	.	.	*	.	.	.
117.3	.	.	.	.	.	.	.	.	.	*	.	.5	.
119.0	.	.	.	.	.	.	.	.	2	*	4	.	.
120.7	.	.	.	.	.	3	1	.	.	*	.	.	.
122.4	.	.	.	.	.	.	.	.	.	*	.	5	.
124.1	.	.	.	.	.	.	.	.	2	*	4	.	.
125.8	.	.	.	.	.	3	1	.	.	*	.	.	.
127.5	.	.	.	.	.	.	.	.	.	*	.	.	.
129.2	.	.	.	.	.	.	.	.	.	*	.	.	.
130.9	.	.	.	.	.	.	.	.	.	*	.	.	.
132.6	.	.	.	.	.	.	.	.	.	*	4	5	.
134.3	.	.	.	.	.	.	1	.	2	*	.	.	.
136.0	.	.	.	.	.	3	.	.	.	*	.	.	.
137.7	.	.	.	.	.	.	.	.	.	*	.	.	.
139.4	.	.	.	.	.	.	.	.	.	*	4	5	.
141.1	.	.	.	.	.	.	.	2	.	*	.	.	.
142.8	.	.	.	3	.	1	.	.	.	*	.	.	.
144.5	.	.	.	.	.	.	.	.	.	*	.	.	.
146.2	.	.	.	.	.	.	.	.	.	*	.	.	.
147.9	.	.	.	.	.	.	.	.	.	*	.	.	.
149.6	.	.	.	.	.	.	.	.	.	*	.	.	.
151.3	.	.	.	.	.	.	.	.	.	*	.	.	.
153.0	.	.	.	.	.	.	.	.	.	*	.	.	.
154.7	.	.	.	.	.	.	.	.	.	*	.	.	.
156.4	.	.	.	.	.	.	.	.	.	*	.	.5	4
158.1	.	.	.	.	.	.	.	.	.	*	.	.	4
159.8	.	.	.	.	.	2	.1	.	.	*	.	5	.
161.5	.	.	.	.3	.	2	.1	.	.	*	.	.	.
163.2	.	.	.	.	3	.	.	.	.	*	.	.	.
164.9	.	.	.	.	.	.	.	.	.	*	.	.	.
166.6	.	.	.	.	.	.	.	.	.	*	.	.	.
168.3	.	.	.	.	.	.	.	.	.	*	.	.	.
170.0	.	.	.	.	.	.	.	.	.	*	.	.	.

MAP SUBAREA SUMMARY FOR FRENCH BROAD, ROSMAN (MAP-0343990 ). PERIOD OF RECORD 10/1959 THRU 09/1961.

WATER YEAR	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	TOTAL
1960	12.80	3.16	5.79	6.84	9.15	7.85	4.30	3.81	6.58	4.37	9.21	6.48	80.33
1961	6.11	1.16	3.36	2.82	11.06	7.05	5.73	4.66	9.60	7.27	17.96	2.30	79.08
	9.46	2.16	4.57	4.83	10.11	7.45	5.01	4.23	8.09	5.82	13.59	4.39	79.71 IN
	WINTER SEASON TOTAL (OCT -APR ) = 43.59 IN												
	SUMMER SEASON TOTAL (MAY -SEP ) = 36.12 IN												

[Back] [Next] [Previous] [Bookmarks] [Top]

Figure 2. Sample output from program MAP

MAP SUBAREA SUMMARY FOR BLANTYRE LOCAL (MAP-0344300 ). PERIOD OF RECORD 10/1959 THRU 09/1961.

WATER YEAR	OCT	NOV	DEC	JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	TOTAL
1960	11.30	2.56	5.20	6.33	9.13	7.57	4.19	3.74	6.92	4.93	9.11	5.18	76.15
1961	6.96	1.06	3.32	2.91	9.47	6.46	5.55	4.91	9.72	8.11	19.00	2.19	79.65
	9.13	1.81	4.26	4.62	9.30	7.01	4.87	4.33	8.32	6.52	14.05	3.68	77.90 IN
													WINTER SEASON TOTAL (OCT -APR ) = 41.00 IN
													SUMMER SEASON TOTAL (MAY -SEP ) = 36.90 IN

\*\*\*\*\*  
COMPLETION CODE = 0  
CPU TIME USED = 0 MINUTES, 9 SECONDS  
CLOCK TIME USED = 0 MINUTES, 18 SECONDS  
\*\*\*\*\*

[Top]