

NOAA Technical Memorandum NWS WR-173

CENTRAL SAN JOAQUIN VALLEY TYPE MAPS

Thomas R. Crossan

Weather Service Office
Fresno, California
December 1981

UNITED STATES
DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary

National Oceanic and
Atmospheric Administration
John V. Byrne, Administrator

National Weather
Service
Richard E. Hallgren, Director



This publication has been reviewed and
is approved for publication by Scientific
Services Division, Western Region.

A handwritten signature in black ink, appearing to read "L. W. Snellman". The signature is written in a cursive style with a long, sweeping tail on the final letter.

L. W. Snellman, Chief
Scientific Services Division
Western Region Headquarters
Salt Lake City, Utah

CONTENTS

	<u>Page</u>
Figures	iv
I. Introduction	1
II. Type Map Classification.....	1
III. Classifying Criteria	3
IV. Examples	4
V. Comparison with Precipitation Type Maps	4

FIGURES

	<u>Page</u>
Figure 1.	Tulare, Kern, and Fresno Counties, California 2
Figure 2.	Type Map 1A-1 6
Figure 3.	Type Map 1A-2 6
Figure 4.	Type Map 1A-3 7
Figure 5.	Type Map 1A-4 7
Figure 6.	Type Map 2A-1 8
Figure 7.	Type Map 2A-2 8
Figure 8.	Type Map 2A-3 9
Figure 9.	Type Map 2B-1 9
Figure 10.	Type Map 2B-2 10
Figure 11.	Type Map 2B-3 10
Figure 12.	Type Map 2C-1 11
Figure 13.	Type Map 2C-2 11
Figure 14.	Type Map 2C-3 12

Central San Joaquin Valley Type Maps

Thomas R. Crossan
Weather Service Office
Fresno, California

I. INTRODUCTION

Type maps have been used as forecast tools by many forecasters over past years. The Fruit-Frost Service in the western states has used type maps effectively for a very long period. There are various methods and procedures to identify or classify maps. This paper is about map typing aimed at a specific forecast problem—forecasting critical minimum temperatures. The various synoptic patterns that precede low minimum temperatures in the Great Valley of California have been typed. Frost or subfreezing temperatures have occurred with all of these types. Of course, the development of damaging temperatures and their duration can be estimated on some occasions without making reference to type maps by just using one or two factors (e.g., low afternoon temperatures, low dew points, cloudy or cloudless skies, etc.). However, type maps can alert one as to whether there may or may not be a damaging-temperature forecast problem when critical temperatures are not obvious. This can be important when preparing agricultural forecasts under operational deadlines.

II. TYPE MAP CLASSIFICATION

Essentially the same type-map classifications are used when preparing forecasts for Tulare, Kern, and Fresno counties (Figure 1). Basic types for the Great Valley of California were originally discussed in Charles C. Allen's paper, "Forecasting Notes on the Central California Citrus District", (about 1954). The subtypes currently in use and described here were added to these basic types by the author.

The classification system is based on using the major upper-air feature; i.e., a ridge or a trough, and its geographical position. Following these first two identifications, the surface map and the lower level windflow determine the final classification. Sea-level isobars over the state of California and its environs are used to describe the surface map. The 700-mb streamline pattern is used to define the lower level wind-flow.

A ridge dominating the west coast is known as Type 1; a trough dominating the same area is classified as Type 2. There are four different surface-map and winds-aloft features used to separate synoptic patterns within the Type 1 classification. Type 1 is usually a stagnant or slow moving weather pattern associated with fog or low clouds for several days or a long string of cold clear "radiational" nights.

Type 2, a trough dominating the west coast, is associated with the majority of cold nights experienced in the central valley. A cold night is defined here as one where the temperature is 32° or lower somewhere in the area. There are three different trough patterns used to separate maps in this type. The first has the trough line just west of the Sierra Nevada; the second has the trough line east of the Sierra Nevada; and the third involves closed lows. Within this third subclassification, the position of the low center itself creates a further separation

for typing. In the first instance the low center is over the Colorado River. In the second the center is off the west coast. In the third the low center is again over the Colorado River, but the upper wind flow over the Great Valley is from the northwest as opposed to southeast in the first instance.

The minimum-temperature forecast problem is most difficult when the arrival of a new air mass behind a frontal system is involved. This is taken into account in the basic classification by appending an "f" to the type map classification to identify the involvement of a frontal passage and/or an air-mass change. Preparing minimum-temperature forecasts during these regimes is obviously more difficult since it involves evaluating the air-mass temperature change in addition to radiational considerations.

III. CLASSIFYING CRITERIA

Type 1 - Ridge over the Pacific coast states

A. Crest overhead

1. NW-SE isobars, with NW winds aloft
2. California trough*, with NE winds aloft
3. California trough*, with W to N winds aloft
4. NE-SW isobars

Type 2 - Trough over the Pacific coast states

A. Trough overhead

1. NW-SE isobars, with NW winds aloft
2. NW-SE isobars, with S-SW winds aloft
3. California trough

B. Trough east of Sierras

1. NW-SE isobars, with W to N winds aloft
2. NE-SW isobars, with N-NE winds aloft
3. California trough, with N winds aloft

C. Closed low (over the Pacific southwest)

1. Center over the Colorado River, SW winds aloft
2. Center off the coast, SW winds aloft
3. Center over the Colorado River, NW winds aloft

Example of these map types are given in Figures 2 through 14. They are used with both forecast and observed flow patterns. Accompanying each type map is:

1. A verbal description of weather, wind-flow, and expected temperatures.
2. The range of dew points usually experienced.

*Lower pressure over the interior sections of the state at surface with higher pressure over the eastern Pacific Ocean and the Nevada Plateau.

3. The applicable range of dry bulb temperatures at 1640 PST.
4. The minimum temperature forecast for the next 18 hours by use of this formula :

$$T_n = D - \frac{H-30}{4} + V + V^I$$

where: D = 1600 PST dew point in F
 H = 1600 PST humidity in %
 V = Variable dependent on D
 V^I = Variable dependent on H

5. The range of station-pressure gradient between Sacramento and Las Vegas at 1600 PST (SAC-LAS). A value of plus 4 is more-or-less normal. Large positive values usually indicate a low probability of frost since they are associated with strong northwesterly winds. Near zero and negative values indicate downslope southeasterly winds (usually involving subsidence and dynamic warming), and a low frost danger.

IV. EXAMPLES

Data for two or three specific, but typical situations, are included with each type map to give the reader a better feel for the potential usefulness of given type maps. Observed late afternoon and evening data given are:

Dry bulb temperatures at 1640 and 1900 PST
Wet bulb temperatures at 1640 and 1900 PST
Max temperature that afternoon
Dew-point temperature (DP) at 1640 and 1900 PST
Relative humidity (RH) at 1640 and 1900 PST
Computed minimum temperature by formula (FORM)
Observed 1600 PST dew points at Sacramento (SAC), Fresno (FNO),
 Bakersfield (BFL), and Coalinga (CLG)*.
Station Pressure gradient (GRAD) between Sacramento and Las Vegas at
 1600 PST.

In the extreme left-hand column of the examples is the minimum temperature that actually occurred the following morning at the stations listed (by initials) in the second column. The stations are all in Tulare County along the Sierra foothills going from north to south: OC-Orange Cove, LC-Lemon Cove, D-Dinuba, I-Ivanhoe, V-Visalia, E-Exeter, T-Tulare, L-Lindsay, P-Porterville, S-Success, TB-Terra Bella, and E-Earlimart. In essence, these temperatures indicate the range of minimum temperatures to be expected with each type.

V. COMPARISON WITH PRECIPITATION TYPE MAPS

The type maps given above were developed as a forecast tool to assist in preparing frost forecasts--mainly on clear and calm nights. The publication of the Map Type Precipitation Probabilities for the western United States, NWS WR-96, were developed to assist forecasters in preparing precipitation forecasts. The

*now closed.

relationship between the frost and precipitation type maps is given in the table below. Column one is the "precipitation" type, column two the corresponding "frost" type, and column three is the probability of precipitation at Fresno.

Winter	1	2B	7%	Summer	1	1A	0
	2	2A	9		2	2B	0
	3	2A	14		3	2A	0
	4	2C3	12		4	2A	0
	5	2C2	10		5	2B	2
	6	1A	1		6	1A	0
	7	2B1	45		7	2B	0
	8	2C2	0		8	1A	2
	9	1A	0		9	2C1	0
	10	2B	50		10	1A	0
Spring	1	2A	14	Fall	1	2B	5
	2	1A	1		2	2B	10
	3	1A	2		3	1A	2
	4	1A	4		4	2B	0
	5	2C3	21		5	1A	4
	6	2C3	4		6	2C1	8
	7	2C3	14		7	2C1	25
	8	2A	33		8	2B	25
	9	2C1	28				
	10	2A	60				

Under 500-mb ridge flow (Frost type 1A) one would not expect much precipitation. Note that this type corresponds to several "precipitation" type maps, and the percentage chance of precipitation under this type is never greater than 4%.

Much of Fresno's annual precipitation occurs in conjunction with upper level closed lows. Under these conditions (Type 2C) precipitation probabilities range from 0 to 28. The other pattern associated with precipitation is with an upper-air trough off the west coast and/or a frontal passage of some sort. These situations are mostly type 2A and occasionally type 2B.

In the writer's opinion the two typing systems correlate well and are synoptically comparable and compatible. Note that in the summer, no matter what the type, there is always a very low chance of precipitation. Climatologically the Great California Valley is dry June through September.


Reference:

Ellison, Eckley S., "A Critique on the Construction and Use of Minimum-Temperature Formulas", Monthly Weather Review, Volume 56, No. 12 (December 1928), pp. 485-495.

Figure 2. Type 1A-1

CREST OVERHEAD DP 31-48
 NW-SE ISOBARS 1640 52-68
 NW WINDS ALOFT FORM 27-30
 GRAD 4-12

THIS TYPE OFTEN HAS CLOUDS, BUT NOT ALWAYS.
 NO EXTREMELY LOW TEMPERATURES OCCUR WITH THIS
 TYPE.




EXAMPLES: 2/19/52 3/17/53

DRY	OC	52/38	38	61/46	35
WET	LC	44/38	37	50/44	36
MAX	D	57	37	72	34
DP	I	34/37	36	40/42	34
RH	V	50/95	38	46/88	35
	E		35		34
	T		36		31
FORM	L	27	35	32	34
SAC	P	43	35	23	35
FNO	S	33	36	37	35
BFL	TB	32	35	36	34
CLG	E	32	37	30	35
GRAD		+4.7		+10.5	

Figure 3. Type 1A-2

CREST OVERHEAD DP 22-40
 CALIF TROUGH 1640 59-70
 NE WINDS ALOFT FORM 25-32
 GRAD <5

THIS IS A CLEAR SKIES VERY LITTLE WIND TYPE EVEN
 THOUGH THE WIND MIGHT BE EXPECTED FROM THE NE
 WINDS ALOFT. THE FORMULA IS RELIABLE EVEN FOR
 LOW RESULTS AND CAN BE COUNTED ON. THE DEW POINT



AT 1900 ORDINARILY SHOULD RISE ABOUT 10 DEGREES
 IF THIS DOES NOT OCCUR EXPECT THE MINIMUM TO GO
 THROUGH THE FORMULA ABOUT 1 DEGREE.


EXAMPLES: 2/21/53 2/25/53 3/14/54

DRY	OC	59/36	28	63/41	33	60/43	31
WET	LC	44/34	30	48/39	32	46/41	30
MAX	D	64	26	67	30	63	32
DP	I	22/32	27	32/37	31	28/38	32
RH	V	23/86	28	32/87	32	29/85	31
	E		25		30		29
	T		26		30		29
FORM	L	25	24	30	29		29
SAC	P	24	26	28	32	23	30
FNO	S	20	27	37	31	32	30
BFL	TB	16	27	33	30	30	29
CLG	E	24	25	27	32	23	30
GRAD		-1.4		+1.7		+1.7	

Figure 4. Type IA-3

CREST OVERHEAD DP 38-45
 CALIF TROUGH 1640 55-63
 W-N WINDS ALOFT FORM 32
 GRAD NEAR 0

THIS IS A HIGH DEW POINT AND FORMULA TYPE AND CLEAR AND CALM CAN BE EXPECTED. FORMULA IS MOST RELIABLE (AS IT IS USUALLY SO HIGH) EXCEPT FOR A 1900 DEW POINT DROP. EVEN SO THE MINIMUMS WILL WILL NOT GO BELOW 30 DEGREES.




EXAMPLES: 2/19/54 3/11/51

DRY	OC	54/40	34	63/43	37
WET	LC	49/39	31	51/42	35
MAX	D	59	32	67	32
DP	I	43/39	34	38/41	34
RH	V	65/96	34	41/92	33
	E		30		32
	T		30		32
FORM	L	32	31	31	32
SAC	P	34	31	MM	34
FNO	S	36	33	MM	32
BFL	TB	37	32	MM	34
CLG	E	36	33	22	34
GRAD		+2.4		-0.4	

Figure 5. Type IA-4

CREST OVERHEAD DP 36
 NE-SW ISOBARS 1640 52-55
 FORM 28-29
 GRAD 0-10

COUNT ON CLEAR SKIES DURING THE NIGHT AND LITTLE WIND. UNUSUALLY CALM WITH GRAD LESS THE 2. WHEN ANTICIPATING CALM CONDITIONS EXPECT MINIMUMS TO GO THROUGH THE FORMULA WITH LC, S, T, AND E GOING BELOW THE L MINIMUM.




EXAMPLES: 2/19/51 2/18/54

DRY	OC	56/28	30	52/48	33
WET	LC	47/38	29	45/45	30
MAX	D	61	29	57	31
DP	I	37/37	30	37/42	32
RH	V	49/46	33	57/83	32
	E		28		30
	T		29		30
FORM	L	29	28	29	31
SAC	P	MM	30	32	30
FNO	S	36	31	37	30
BFL	TB	27	29	35	32
CLG	E	28	28	24	32
GRAD		+1.3		+10.1	

Figure 6. Type 2A-1

TROUGH OVERHEAD DP 24-41
 NW-SE ISOBARS 1640 46-61
 NW WINDS ALOFT FORM 22-32
 GRAD 4-16
 (MOST >10)




USUALLY A WINDY TYPE WITH CLEAR SKIES. SOME REALLY GET COLD. CONSIDER THE FORMULA RELIABLE WITH 2-3 DEGREES ADDED ON WINDY NIGHTS. AFTER A FRONTAL PASSAGE EXPECT THE FORMULA TO BE TOO HIGH. IN GENERAL THE LOWER DP AND FORM ARE THE COLDER ONES, BUT CLOUDS ARE NOT TO BE COUNTED OUT. THIS TYPE IS FULL OF 'TRICKY' ONES- BE CAREFUL.

EXAMPLES:		4/8/53		2/17/52		3/10/54	
DRY	OC	57/43	29	49/41	29	51/46	31
WET	LC	44/38	28	43/39	28	46/43	30
MAX	D	63	28	60	MM	61	33
DP	I	24/32	30	37/37	28	42/40	32
RH	V	28/65	29	62/86	MM	70/77	31
	E		29		27	30	31
	T		28		27	29	33
FORM	L	25	29	28	26	30	31
SAC	P	29	26	24	28	34	30
FNO	S	23	26	27	27	37	31
BFL	TB	23	27	42	28	46	32
CLG	E	30	32	38	29	32	32
GRAD		+9.1		+12.2		+16.6	

Figure 7. Type 2A-2

TROUGH OVERHEAD DP 30-38
 NW-SE ISOBARS 1640 52-70
 S-SW WINDS ALOFT FORM 28-33
 GRAD 2 OR 9
 (NO IN-BETWEEN)




USUALLY CLEAR SKIES WITH THIS TYPE AND THE FORMULA IS QUITE RELIABLE EXCEPT IN THE 1900 DP DROP IF THE 1900 TMP IS >45 THE LOWEST MIN WILL BE 31 UNLESS THERE IS A DP DROP- THEN 30. WITH THE 1900 TMP <45 THE MIN WILL GO AS LOW AS 28. WATCH FOR TULARE TO GO BELOW THE LINDSAY KEY STATION MINIMUM. THE RULE APPLIES ASSUMING NO FRONTAL PASSAGE.

EXAMPLES:		2/20/51		2/21/51		3/13/53	
DRY	OC	61/40	32	52/47	29	65/45	36
WET	LC	48/39	29	48/43	30	51/44	35
MAX	D	66	33	63	31	69	37
DP	I	33/38	32	41/39	30	37/42	35
RH	V	35/92	32	66/73	30	35/90	34
	E		32		30		33
	T		28		29		34
FORM	L	30	31	31	30	33	33
SAC	P	46	30	40	33	33	34
FNO	S	35	30	39	34	MM	34
BFL	TB	22	33	34	31	35	34
CLG	E	30	30	36	31	33	34
GRAD		+1.7		+9.5		+1.1	

Figure 8. Type 2A-3

TROUGH OVERHEAD DP 40-45
 CALIF TROUGH 1640 55-65
 FORM NR 32
 GRAD 2-9

THIS IS A CLEAR SKIES TYPE WITH HIGH DEW POINT AND FORMULA. THIS TYPE IS NEVER TOO COLD AND HAS YET TO HAVE A TEMPERATURE BELOW 30.




EXAMPLES:

		3/4/55		3/5/55	
DRY	OC	60/44	36	60/44	36
WET	LC	51/43	35	50/42	34
MAX	D	65	33	63	35
DP	I	43/42	34	41/41	35
RH	V	54/93	32	49/92	35
	E		33		32
	T		30		34
FORM	L	32	33	31	33
SAC	P	22	34	19	36
FNO	S	40	37	40	34
BFL	TB	37	34	37	33
CLG	E	39	32	29	33
GRAD		+7.1		+2.0	

Figure 9. Type 2B-1

PLATEAU TROUGH DP 23-44
 NW-SE ISOBARS 1640 50-66
 W-N WINDS ALOFT FORM 24-33
 GRAD 5-18
 (MOST 8-12)

THIS TYPE INCLUDES ALL FORECAST PROBLEMS OF WIND CLOUDS AND A WIDE RANGE OF DEW POINT. DISREGARDING APRIL TYPES, A DP BELOW 30 USUALLY GIVES MINS BELOW 30. WIND CAN USUALLY BE COUNTED ON AND 3 DEGREES ADDED TO THE FORMULA WHEN IT IS



BELOW 27, BUT USE CAUTION IN ADDING TO HIGHER FORMULAS. CLOUDS AND WIND WITH THIS TYPE GIVES A WIDE RANGE IN MINS THROUGHOUT THE AREA. WITH THE WINDY TYPES, BE CAREFUL OF S AND LC GOING 2-3 BELOW THE KEY STATION MINIMUM.


EXAMPLES:

		3/21/53		3/11/54		4/10/53
DRY	OC	50/34	30	50/42	31	59/50
WET	LC	42/34	31	41/39	28	45/41
MAX	D	55	30	55	34	63
DP	I	31/33	29	30/36	30	26/29
RH	V	49/45	29	46/80	30	28/44
	E		29		29	35
	T		29		33	31
FORM	L	24	28	24	28	26
SAC	P	20	28	33	28	36
FNO	S	27	27	29	30	32
BFL	TB	31	27	27	28	27
CLG	E	18	28	35	32	29
GRAD		+7.1		+11.5		+13.2

Figure 10. Type 2B-2

PLATEAU TROUGH DP 24-32
 NE-SW ISOBARS 1640 51-60
 N-NE WINDS ALOFT FORM 23-27
 GRAD 3-10
 (MOST NR 10)

COUNT ON CLEAR SKIES AND OCCASSIONAL BREEZES. LOW VALLEY DEW POINTS ARE THE RULE AND THE FORMULA IS USUALLY 2 DEGREES LOW DEPENDING ON THE WIND. EVEN THE COLDEST HAD A FORM 2 DEGREES BELOW THE MIN. SOME STATIONS WILL GO AS LOW AS THE FORMULA EVEN THOUGH NOT AT THE KEY STATION. WITH THIS WINDY TYPE, BE CAREFUL OF S, E, T, AND LC GOING 2-3 BELOW THE KEY STATION. EXPECT THE BOTTOM TO DROP OUT WITH FRONTAL PASSAGE.




EXAMPLES:	2/19/53	2/18/53	3/20/52
DRY OC	51/34 29	53/36 27	54/46 31
WET LC	40/32 30	43/36 26	44/43 30
MAX D	56 26	55 27	57 31
DP I	24/31 29	30/35 28	32/40 30
RH V	35/90 28	43/94 30	43/78 34
E		25	26
T		25	24
FORM L	23 26 25 28 27 31		
SAC P	15 28 20 29 30 31		
FNO S	23 25 28 25 34 29		
BFL TB	27 28 31 27 31 31		
CLG E	16 25 9 23 22 31		
GRAD	+9.1	+12.5	+10.1

Figure 11. Type 2B-3

PLATEAU TROUGH DP 23-32
 CALIF TROUGH 1640 55-60
 N WINDS ALOFT FORM 25-30
 GRAD 1-7
 (MOST NR 0)

ON THIS TYPE THE DEW POINT AND FORMULA INDICATE THE WARM AND COLD ONES. DP BELOW 25 GIVES 22-28 DP 25-32 GIVES 28-32; AND DP NEAR 32 GIVES NEAR 32. WIND AND CLOUDS OCCASSIONALLY A PROBLEM, BUT USUALLY CLEAR SKIES CAN BE COUNTED ON AND THE FORMULA CONSIDERED RELIABLE. IN MARCH AND APRIL WHEN THE DP RISES BY 10 OR MORE- ADD 4 TO THE FORMULA FOR THE KEY STATION MINIMUM.




EXAMPLES:	2/20/53	3/3/53	3/22/52
DRY OC	56/36 28	58/40 31	58/40 35
WET LC	42/33 32	47/38 30	45/39 34
MAX D	60 24	61 30	62 35
DP I	23/28 26	34/37 32	29/28 34
RH V	28/73 28	40/89 32	34/93 33
E		24	29
T		24	29
FORM L	25 22 30 29 26 31		
SAC P	21 25 33 30 30 33		
FNO S	22 25 28 31 33 31		
BFL TB	11 24 33 29 31 33		
CLG E	18 22 26 31 33 32		
GRAD	+1.1	+1.8	+2.7

Figure 12. Type 2C-1

CLOSED LOW DP 37
 COLORADO RIVER 1640 49
 FORM 28
 GRAD 6.

LOW VALLEY DEW POINTS ARE THE RULE WITH NE WINDS ALOFT. FORMULA RELIABLE UNLESS 1900 DEW POINTS- THEN EXPECT MINIMUM BELOW THE FORMULA. WIND IS THE PROBLEM HERE RATHER THAN CLOUDS.




EXAMPLES: 2/23/53

DRY	OC	48/35	28
WET	LC	43/34	27
MAX	D	55	26
DP	I	36/33	28
RH	V	63	29
	E		26
	T		26
FORM	L	28	27
SAC	P	13	27
FNO	S	37	26
BFL	TB	37	27
CLG	E	16	28
GRAD		+5.8	

Figure 13. Type 2C-2

CLOSED LOW DP 27-47
 CNTR OFF COAST 1640 45-61
 FORM 24-32
 GRAD 0-9
 (MOST 3)

INVARIABLY THE 700 MB TEMPS ARE -8 OR LOWER. CLOUDINESS IS A CAUSE FOR CONCERN CAUSING ABOVE 32 ON LOW FORMULAS. WITH CLEAR SKIES, FORMULA WILL BE CORRECT UNLESS THE DP DROPS AT 1900, THEN IT WILL BE TOO HIGH. BIG PROBLEM IS THE EFFECT OF CLOUDS, IF ANY; NOT WIND.




EXAMPLES: 2/23/51 3/7/52 3/16/54

DRY	OC	47/45	30	46/36	31	41/41	38
WET	LC	45/42	30	43/36	32	41/40	39
MAX	D	62	29	51	30	62	38
DP	I	43/38	30	40/35	30	40/40	38
RH	V	87/76	30	81/97	30	78/97	38
	E		29		29		38
	T		27		30		36
FORM	L	29	28	27	28	28	36
SAC	P	35	30	MM	30	41	40
FNO	S	37	29	39	28	43	37
BFL	TB	30	29	40	29	40	37
CLG	E	41	28	37	31	40	38
GRAD		+0.0		+2.0		+3.4	

Figure 14. Type 2C-3

CLOSED LOW DP 29-43
 COLORADO RIVER 1640 46-67
 FORM 22-31
 GRAD 4-10

WITH A 22-26 FORMULA AND DP LESS THAN 29 THE MINIMUMS WILL RANGE 25-28. WITH FORM 27-31 AND DP >30, MINS WILL BE 29 AND HIGHER. HIGHER FORM AND DP GIVES ABOVE 32 DUE TO CLOUDS FORMED IN THE COLD AIR ALOFT. COLD AIR ALOFT DOES NOT NECESARILLY MEAN CLOUDS HOWEVER. WITH A FORM OF 22-24 EXPECT SOME STATIONS TO GO THAT LOW, BUT PERHAPS NOT THE LINDSAY KEY STATIONS.



EXAMPLES:		3/2/51		3/1/53		3/31/51	
DRY	OC	46/37	26	47/38	27	67/47	36
WET	LC	39/36	29	40/38	26	50/43	42
MAX	D	53	27	52	30	70	32
DP	I	29/35	28	30/36	29	30/39	35
RH	V	50/91	28	50/91	28	25/72	37
	E		25		28		31
	T		26		28		33
FORM	L	22	26	23	27	27	32
SAC	P	20	27	29	27	30	35
FNO	S	29	27	35	25	32	35
BFL	TB	31	27	35	27	18	32
CLG	E	24	31	28	30	24	39
GRAD		+8.8		+6.8		+4.7	

- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Vulcan Flat Weather Station. R. F. Quiring, June 1977. (PB-271-764/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Webb, Jr., June 1977. (PB-271-742/AS)
- 123 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Lefterich, August 1977. (PB-272-661)
- 124 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Lefterich and Charles J. Neumann, August 1977. (PB-273-155/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Mielke, February 1978. (PB-281-387/AS)
- 128 Band Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudgal, April 1978. (PB-283-880/AS)
- 129 Fire Warnings. David W. Coans, May 1978. (PB-283-866/AS)
- 130 Flood-Flood Procedure. Ralph E. Hatch and Gerald Williams, May 1978. (PB-286-014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Holliner and David E. Olson, September 1978. (PB-289-916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-746 Weather Radar. R. G. Pappas, R. V. Lee, D. W. Finke, October 1978. (PB289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., Jan. 1979. (PB294324/AS)
- 138 A Simple Analytic/Diagnostic System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)
- 139 Aids for Forecasting Minimum Temperature in the Mendocino Frost District. Robert S. Robinson, April 1979. (PB298359/AS)
- 140 Influence of Cloudiness on Summer-time Temperatures in the Eastern Washington Fire Weather District. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and WRF Precipitation Guidance for Nevada During Uvran. Christopher Hill, April 1979. (PB298619/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Gorey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 145 On the Use of Solar Radiation and Temperature Models to Estimate the Snap Beam Maturity Date in the Willamette Valley. Earl W. Bates, August 1979. (PB80-160971)
- 146 The BARE Experiment. Morris S. Webb, October 1979. (PB80-155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80-160344)
- 148 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80-174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80-226485)
- 151 WRF Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80-196039)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins, June 1980. (PB80-225493) (Out of Print.)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80-225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Gard, July 1980. (PB81-108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Tenharkel, July 1980. (PB81-106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81-102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Cohen, August 1980. (PB81-108227)
- 158 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81-109779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Lefterich and Emil M. Brown, February 1981.
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Barry Anderson, April 1981.
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J. G. Burnash and R. Larry Ferral, April 1981.
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. Lt. Colonel Keith R. Goodley, April 1981.
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981.
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981.
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981.
- 167 ARAP User's Guide. Mark Mathewson, July 1981. (revised September 1981)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981.
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Lefterich, Jr., October 1981.
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981.
- 171 Verification of 72-hour 500-mb Height Predictions. R. F. Quiring, November 1981.
- 172 Forecasting Heavy Snow at Mount Rainier, Washington. James W. Holcomb, December 1981.

NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

The National Oceanic and Atmospheric Administration was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

PROFESSIONAL PAPERS — Important definitive research results, major techniques, and special investigations.

CONTRACT AND GRANT REPORTS — Reports prepared by contractors or grantees under NOAA sponsorship.

ATLAS — Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

TECHNICAL SERVICE PUBLICATIONS — Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

TECHNICAL REPORTS — Journal quality with extensive details, mathematical developments, or data listings.

TECHNICAL MEMORANDUMS — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



Information on availability of NOAA publications can be obtained from:

**ENVIRONMENTAL SCIENCE INFORMATION CENTER (D822)
ENVIRONMENTAL DATA AND INFORMATION SERVICE
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION
U.S. DEPARTMENT OF COMMERCE**

**6009 Executive Boulevard
Rockville, MD 20852**