

NOAA Technical Memorandum NWS WR-198



---

PRELIMINARY LIGHTNING CLIMATOLOGY STUDIES FOR IDAHO

Christopher D. Hill  
Carl J. Gorski  
Michael C. Conger

Salt Lake City, Utah  
April 1987

---

**U.S. DEPARTMENT OF  
COMMERCE**

National Oceanic and  
Atmospheric Administration

National Weather  
Service

4  
M94.221:82.2  
Hil  
1987



NOAA TECHNICAL MEMORANDA  
National Weather Service, Western Region Subseries

The National Weather Service (NWS) Western Region (WR) Subseries provides an informal medium for the documentation and quick dissemination of results not appropriate, or not yet ready, for formal publication. The series is used to report on work in progress, to describe technical procedures and practices, or to relate progress to a limited audience. These Technical Memoranda will report on investigations devoted primarily to regional and local problems of interest mainly to personnel, and hence will not be widely distributed.

Papers 1 to 25 are in the former series, ESSA Technical Memoranda, Western Region Technical Memoranda (WRTM); papers 24 to 59 are in the former series, ESSA Technical Memoranda, Weather Bureau Technical Memoranda (NBTM). Beginning with 60, the papers are part of the series, NOAA Technical Memoranda NWS. Out-of-print memoranda are not listed.

Papers 2 to 22, except for 5 (revised edition), are available from the National Weather Service Western Region, Scientific Services Division, P.O. Box 11188, Federal Building, 125 South State Street, Salt Lake City, Utah 84147. Paper 5 (revised edition), and all others beginning with 25 are available from the National Technical Information Service, U.S. Department of Commerce, Sills Building, 5285 Port Royal Road, Springfield, Virginia 22161. Prices vary for all paper copies; \$3.50 microfiche. Order by accession number shown in parentheses at end of each entry.

ESSA Technical Memoranda (WRTM)

- 2 Climatological Precipitation Probabilities. Compiled by Lucianne Miller, December 1965.
- 3 Western Region Pre- and Post-FP-3 Program, December 1, 1965, to February 20, 1966. Edward D. Diemer, March 1966.
- 5 Station Descriptions of Local Effects on Synoptic Weather Patterns. Philip Williams, Jr., April 1966 (revised November 1967, October 1969). (PB-17800)
- 8 Interpreting the RAREP. Herbert P. Benner, May 1966 (revised January 1967).
- 11 Some Electrical Processes in the Atmosphere. J. Latham, June 1966.
- 17 A Digitalized Summary of Radar Echoes within 100 Miles of Sacramento, California. J. A. Youngberg and L. B. Overaas, December 1966.
- 21 An Objective Aid for Forecasting the End of East Winds in the Columbia Gorge, July through October. D. John Coparanis, April 1967.
- 22 Derivation of Radar Horizons in Mountainous Terrain. Roger G. Pappas, April 1967.

ESSA Technical Memoranda, Weather Bureau Technical Memoranda (WBTM)

- 25 Verification of Operation Probability of Precipitation Forecasts, April 1966-March 1967. W. W. Dickey, October 1967. (PB-176240)
- 26 A Study of Winds in the Lake Mead Recreation Area. R. P. Augulis, January 1968. (PB-177830)
- 28 Weather Extremes. R. J. Schmidl, April 1968 (revised March 1986). (PB86 17672/AS)
- 29 Small-Scale Analysis and Prediction. Philip Williams, Jr., May 1968. (PB178425)
- 30 Numerical Weather Prediction and Synoptic Meteorology. CPT Thomas D. Murphy, USAF, May 1968. (AD 673365)
- 31 Precipitation Detection Probabilities by Salt Lake ARTC Radars. Robert K. Belesky, July 1968. (PB 179084)
- 32 Probability Forecasting--A Problem Analysis with Reference to the Portland Fire Weather District. Harold S. Ayer, July 1968. (PB 179289)
- 36 Temperature Trends in Sacramento--Another Heat Island. Anthony D. Lentini, February 1969. (PB 183055)
- 37 Disposal of Logging Residues Without Damage to Air Quality. Owen P. Cramer, March 1969. (PB 183057)
- 39 Upper-Air Lows Over Northwestern United States. A.L. Jacobson, April 1969. PB 184296)
- 40 The Man-Machine Mix in Applied Weather Forecasting in the 1970s. L.W. Snellman, August 1969. (PB 185066)
- 43 Forecasting Maximum Temperatures at Helena, Montana. David E. Olsen, October 1969. (PB 185762)
- 44 Estimated Return Periods for Short-Duration Precipitation in Arizona. Paul C. Kangieser, October 1969. (PB 187763)
- 46 Applications of the Net Radiometer to Short-Range Fog and Stratus Forecasting at Eugene, Oregon. L. Yee and E. Bates, December 1969. (PB 190476)
- 47 Statistical Analysis as a Flood Routing Tool. Robert J.C. Burnash, December 1969. (PB 188744)
- 48 Tsunami. Richard P. Augulis, February 1970. (PB 190157)
- 49 Predicting Precipitation Type. Robert J.C. Burnash and Floyd E. Hug, March 1970. (PB 190962)
- 50 Statistical Report on Aeroallergens (Pollens and Molds) Fort Huachuca, Arizona, 1969. Wayne S. Johnson, April 1970. (PB 191743)
- 51 Western Region Sea State and Surf Forecaster's Manual. Gordon C. Shields and Gerald B. Burdwell, July 1970. (PB 193102)
- 52 Sacramento Weather Radar Climatology. R.G. Pappas and C. M. Veliquette, July 1970. (PB 193347)
- 54 A Refinement of the Vorticity Field to Delineate Areas of Significant Precipitation. Barry B. Aronovitch, August 1970.
- 55 Application of the SSARR Model to a Basin without Discharge Record. Vail Schermerhorn and Donal W. Kuehl, August 1970. (PB 194394)
- 56 Area Coverage of Precipitation in Northwestern Utah. Philip Williams, Jr., and Werner J. Heck, September 1970. (PB 194389)
- 57 Preliminary Report on Agricultural Field Burning vs. Atmospheric Visibility in the Willamette Valley of Oregon. Earl M. Bates and David O. Chilcote, September 1970. (PB 194710)
- 58 Air Pollution by Jet Aircraft at Seattle-Tacoma Airport. Wallace R. Donaldson, October 1970. (COM 71 00017)
- 59 Application of PE Model Forecast Parameters to Local-Area Forecasting. Leonard W. Snellman, October 1970. (COM 71 00016)

NOAA Technical Memoranda (NWS WR)

- 60 An Aid for Forecasting the Minimum Temperature at Medford, Oregon, Arthur W. Fritz, October 1970. (COM 71 00120)
- 63 700-mb Warm Air Advection as a Forecasting Tool for Montana and Northern Idaho. Norris E. Noerner, February 1971. (COM 71 00349)
- 64 Wind and Weather Regimes at Great Falls, Montana. Warren B. Price, March 1971.
- 66 A Preliminary Report on Correlation of ARTCC Radar Echoes and Precipitation. Wilbur K. Hall, June 1971. (COM 71 00829)

- 69 National Weather Service Support to Soaring Activities. Ellis Burton, August 1971. (COM 71 00956)
- 71 Western Region Synoptic Analysis-Problems and Methods. Philip Williams, Jr., February 1972. (COM 72 10433)
- 74 Thunderstorms and Hail Days Probabilities in Nevada. Clarence M. Sakamoto, April 1972. (COM 72 10554)
- 75 A Study of the Low Level Jet Stream of the San Joaquin Valley. Ronald A. Willis and Philip Williams, Jr., May 1972. (COM 72 10707)
- 76 Monthly Climatological charts of the Behavior of Fog and Low Stratus at Los Angeles International Airport. Donald M. Gales, July 1972. (COM 72 11140)
- 77 A Study of Radar Echo Distribution in Arizona During July and August. John E. Hales, Jr., July 1972. (COM 72 11136)
- 78 Forecasting Precipitation at Bakersfield, California, Using Pressure Gradient Vectors. Earl T. Riddiough, July 1972. (COM 72 11146)
- 79 Climate of Stockton, California. Robert C. Nelson, July 1972. (COM 72 10920)
- 80 Estimation of Number of Days Above or Below Selected Temperatures. Clarence M. Sakamoto, October 1972. (COM 72 10021)
- 81 An Aid for Forecasting Summer Maximum Temperatures at Seattle, Washington. Edgar G. Johnson, November 1972. (COM 73 10150)
- 82 Flash Flood Forecasting and Warning Program in the Western Region. Philip Williams, Jr., Chester L. Glenn, and Roland L. Raetz, December 1972, (revised March 1978). (COM 73 10251)
- 83 A comparison of Manual and Semiautomatic Methods of Digitizing Analog Wind Records. Glenn E. Rasch, March 1973. (COM 73 10669)
- 86 Conditional Probabilities for Sequences of Wet Days at Phoenix, Arizona. Paul C. Kangieser, June 1973. (COM 73 11264)
- 87 A Refinement of the Use of K-Values in Forecasting Thunderstorms in Washington and Oregon. Robert Y.G. Lee, June 1973. (COM 73 11276)
- 89 Objective Forecast Precipitation Over the Western Region of the United States. Julia N. Paegle and Larry P. Kierulff, September 1973. (COM 73 11946/3AS)
- 91 Arizona "Eddy" Tornadoes. Robert S. Ingram, October 1973. (COM 73 10465)
- 92 Smoke Management in the Willamette Valley. Earl M. Bates, May 1974. (COM 74 11277/AS)
- 93 An Operational Evaluation of 500-mb Type Regression Equations. Alexander E. MacDonald, June 1974. (COM 74 11407/AS)
- 94 Conditional Probability of Visibility Less than One-Half Mile in Radiation Fog at Fresno, California. John D. Thomas, August 1974. (COM 74 11555/AS)
- 95 Climate of Flagstaff, Arizona. Paul W. Sorenson, and updated by Reginald W. Preston, January 1987. (PB87 143160/AS)
- 96 Map type Precipitation Probabilities for the Western Region. Glenn E. Rasch and Alexander E. MacDonald, February 1975. (COM 75 10428/AS)
- 97 Eastern Pacific Cut-Off Low of April 21-28, 1974. William J. Alder and George R. Miller, January 1976. (PB 250 711/AS)
- 98 Study on a Significant Precipitation Episode in Western United States. Ira S. Brenner, April 1976. (COM 75 10719/AS)
- 99 A Study of Flash Flood Susceptibility-A Basin in Southern Arizona. Gerald Williams, August 1975. (COM 75 11360/AS)
- 102 A Set of Rules for Forecasting Temperatures in Napa and Sonoma Counties. Wesley L. Tuft, October 1975. (PB 246 902/AS)
- 103 Application of the National Weather Service Flash-Flood Program in the Western Region. Gerald Williams, January 1976. (PB 253 053/AS)
- 104 Objective Aids for Forecasting Minimum Temperatures at Reno, Nevada, During the Summer Months. Christopher D. Hill, January 1976. (PB 252 866/AS)
- 105 Forecasting the Mono Wind. Charles P. Ruscha, Jr., February 1976. (PB 254 650)
- 106 Use of MOS Forecast Parameters in Temperature Forecasting. John C. Plankinton, Jr., March 1976. (PB 254 649)
- 107 Map Types as Aids in Using MOS PoPs in Western United States. Ira S. Brenner, August 1976. (PB 259 594)
- 108 Other Kinds of Wind Shear. Christopher D. Hill, August 1976. (PB 260 437/AS)
- 109 Forecasting North Winds in the Upper Sacramento Valley and Adjoining Forests. Christopher E. Fontana, September 1976. (PB 273 677/AS)
- 110 Cool Inflow as a Weakening Influence on Eastern Pacific Tropical Cyclones. William J. Denney, November 1976. (PB 264 655/AS)
- 112 The MAW/MOS Program. Alexander E. MacDonald, February 1977. (PB 265 941/AS)
- 113 Winter Season Minimum Temperature Formula for Bakersfield, California, Using Multiple Regression. Michael J. Oard, February 1977. (PB 273 694/AS)
- 114 Tropical Cyclone Kathleen. James R. Fors, February 1977. (PB 273 676/AS)
- 116 A Study of Wind Gusts on Lake Mead. Bradley Colman, April 1977. (PB 268 847)
- 117 The Relative Frequency of Cumulonimbus Clouds at the Nevada Test Site as a Function of K-Value. R.F. Quiring, April 1977. (PB 272 831)
- 118 Moisture Distribution Modification by Upward Vertical Motion. Ira S. Brenner, April 1977. (PB 268 740)
- 119 Relative Frequency of Occurrence of Warm Season Echo Activity as a Function of Stability Indices Computed from the Yucca Flat, Nevada, Rawinsonde. Darryl Randerson, June 1977. (PB 271 290/AS)
- 121 Climatological Prediction of Cumulonimbus Clouds in the Vicinity of the Yucca Flat Weather Station. R.F. Quiring, June 1977. (PB 271 704/AS)
- 122 A Method for Transforming Temperature Distribution to Normality. Morris S. Hebb, Jr., June 1977. (PB 271 742/AS)
- 124 Statistical Guidance for Prediction of Eastern North Pacific Tropical Cyclone Motion - Part I. Charles J. Neumann and Preston W. Leftwich, August 1977. (PB 272 661)
- 125 Statistical Guidance on the Prediction of Eastern North Pacific Tropical Cyclone Motion - Part II. Preston W. Leftwich and Charles J. Neumann, August 1977. (PB 273 155/AS)
- 127 Development of a Probability Equation for Winter-Type Precipitation Patterns in Great Falls, Montana. Kenneth B. Hielke, February 1978. (PB 281 387/AS)
- 128 Hand Calculator Program to Compute Parcel Thermal Dynamics. Dan Gudge, April 1978. (PB 283 080/AS)
- 129 Fire whirls. David N. Goens, May 1978. (PB 283 866/AS)
- 130 Flash-Flood Procedure. Ralph C. Hatch and Gerald Williams, May 1978. (PB 286 014/AS)
- 131 Automated Fire-Weather Forecasts. Mark A. Mollner and David E. Olsen, September 1978. (PB 289 916/AS)
- 132 Estimates of the Effects of Terrain Blocking on the Los Angeles WSR-74C Weather Radar. R.G. Pappas, R.Y. Lee, B.W. Finke, October 1978. (PB 289767/AS)
- 133 Spectral Techniques in Ocean Wave Forecasting. John A. Jannuzzi, October 1978. (PB291317/AS)
- 134 Solar Radiation. John A. Jannuzzi, November 1978. (PB291195/AS)
- 135 Application of a Spectrum Analyzer in Forecasting Ocean Swell in Southern California Coastal Waters. Lawrence P. Kierulff, January 1979. (PB292716/AS)
- 136 Basic Hydrologic Principles. Thomas L. Dietrich, January 1979. (PB292247/AS)
- 137 LFM 24-Hour Prediction of Eastern Pacific Cyclones Refined by Satellite Images. John R. Zimmerman and Charles P. Ruscha, Jr., January 1979. (PB294324/AS)
- 138 A Simple Analysis/Diagnosis System for Real Time Evaluation of Vertical Motion. Scott Heflick and James R. Fors, February 1979. (PB294216/AS)

NOAA Technical Memorandum NWS WR-198

PRELIMINARY LIGHTNING CLIMATOLOGY STUDIES

Christopher D. Hill  
Carl J. Gorski  
Michael C. Conger

Weather Service Forecast Office  
National Weather Service  
Boise, Idaho  
April 1987


UNITED STATES  
DEPARTMENT OF COMMERCE  
Malcolm Baldrige, Secretary

National Oceanic and  
Atmospheric Administration  
John V. Byrne, Administrator

National Weather  
Service  
Richard E. Hallgren, Director



This technical publication has been  
reviewed and is approved for  
publication by Scientific Services  
Division, Western Region.

A handwritten signature in black ink that reads "Glenn E. Rasch". The signature is written in a cursive style with a large initial 'G'.

Glenn E. Rasch, Chief  
Scientific Services Division  
Western Region Headquarters  
Salt Lake City, Utah

## TABLE OF CONTENTS

	<u>PAGE</u>
List of Tables	iv
List of Figures	v
Abstract	1
I. Introduction	2
II. Preliminary Thunderstorm Climatology	2
III. Frequency of Lightning Under Various Synoptic Flow Patterns	4
IV. Manually Observed Lightning Activity Level	6
V. Discussion	8
VI. References	9

## LIST OF TABLES

	<u>PAGE</u>
Table 1. Frequency of Correct Categories Verified Against NFDRS Cloud-to-Ground Lightning Density Limits	34
Table 2. Lightning Density Mean, Standard Deviation and Range for Man-Observed Thunderstorm Days by LAL Category	35
Table 3. Number of Cloud-to-Ground Strikes Within 2500 mi <sup>2</sup> Area Per LAL Category	36

## LIST OF FIGURES

		<u>PAGE</u>
Figure 1.	Average Annual Number of Days with Thunderstorms (from <u>Climate and Man</u> , 1941)	11
Figure 2.	Average Annual Number of Days with Thunderstorms (from Critchfield, 1966)	12
Figure 3.	Average Annual Number of Days with Thunderstorms (from Changery, 1981)	13
Figure 4.	Grid Block (One-half by One-half Degree) Array for which Lightning Data Archived and Analyzed	14
Figure 5.	Average Annual Number of Days with Lightning Strikes Based on Two Years (1985, 1986) of Data	15
Figure 6.	Number of Days with Lightning Strikes during 1985	16
Figure 7.	Number of Days with Lightning Strikes during 1986	17
Figure 8.	Average Annual Number of Hours with Lightning Strikes Based on Two Years (1985, 1986) of Data	18
Figure 9.	Average Annual Number of Lightning Strikes Based on Two Years of Data (1985, 1986)	19
Figure 10.	Average Annual Percent of Summer (June, July, August) Days with Lightning Strikes	20
Figure 11.	Summer Map Type 1 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	21
Figure 12.	Summer Map Type 2 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	22
Figure 13.	Summer Map Type 3 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	23
Figure 14.	Summer Map Type 4 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	24
Figure 15.	Summer Map Type 5 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	25

	<u>PAGE</u>
Figure 16. Summer Map Type 6 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	26
Figure 17. Summer Map Type 7 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	27
Figure 18. Summer Map Type 8 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	28
Figure 19. Summer Map Type 9 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	29
Figure 20. Summer Map Type 10 and Corresponding Percent Probabilities of Lightning During the Period Noon to Midnight MDT	30
Figure 21. Map of Verification Stations	31
Figure 22. NFDRS Relative Frequency Distribution vs. Observed Frequency Distribution	32
Figure 23. Frequency Distribution of Correct Observations Using NFDRS Limits vs. Correct Observations Using Adjusted Limits	33

## PRELIMINARY LIGHTNING CLIMATOLOGY STUDIES FOR IDAHO

### ABSTRACT

Two years of cloud-to-ground lightning strike data from the Bureau of Land Management's Automatic Lightning Detection System are used to develop a preliminary thunderstorm climatology for the State of Idaho. The preliminary results suggest earlier climatologies based on widely separated weather observing stations underestimate the number of thunderstorm days in the region. Map types are then used to investigate the variations in the frequency of lightning under differing mid-tropospheric flow patterns. Finally, manually reported lightning activity levels are compared to lightning occurrence and density as measured by the automated system.

## PRELIMINARY LIGHTNING CLIMATOLOGY STUDIES FOR IDAHO

### I. INTRODUCTION

In 1983, the National Weather Service (NWS) Western Region began utilizing cloud-to-ground lightning strike information from the Bureau of Land Management's (BLM) detection network. Since the original description by Rasch and Mathewson (1984), significant changes have been made to both the NWS and BLM programs. Most notably, the BLM has increased the number of sensors, added the capability to differentiate between positive and negative strikes, and extended the program to year-round operation. The NWS now processes and archives the lightning data at the NWS Forecast Office in Boise, Idaho, since the facility is co-located with the BLM computers. The programs of these two agencies came to maturity in 1985. As a result, the data sets for 1985 and 1986 are almost 100 percent complete. These two years of data were used to develop a preliminary thunderstorm climatology for the Idaho area, investigate lightning occurrence under various mid-tropospheric flow patterns, and compare BLM and U.S. Forest Service (USFS) manually reported lightning activity levels (LAL's) with actual lightning occurrence.

### II. PRELIMINARY THUNDERSTORM CLIMATOLOGY

Until now, thunderstorm climatologies have been based on the only available source, surface weather observations. Unfortunately, the western United States in general and Idaho in particular have few surface weather observing sites. In addition, the region is largely mountainous, and the few observation sites that do exist are in valleys. This has led to what appears to be a significant underestimation of the frequency of thunderstorm activity.

An early example of thunderstorm climatology is shown in Figure 1 from *Climate and Man* (1941). Based on 40 years of data the climatology depicts the average annual number of thunderstorm days varying from somewhat less than 20 days in western Idaho to a little more than 40 days per year in eastern Idaho. Figure 2, from Critchfield (1966), portrays a significantly different climatology, with the area in eastern Idaho which was a maximum in Figure 1, now depicted as a relative minimum. A maximum is analyzed over southwest Montana. Figure 3 from Changery (1981) is an attempt to utilize all available data and subjectively incorporate major terrain influences. This climatology depicts a relative minimum through the Snake River Valley of southern Idaho with maxima near the central Idaho-southwest Montana border and just east of Idaho in western Wyoming. A secondary maximum is also shown over the mountains along the Idaho-Utah border.

Since 1983, lightning strike data has been available from the BLM detection network and is transmitted to NWS computers at Boise in real time. Each strike is stored with its time of occurrence, latitude, longitude, signal strength and sign (positive or negative), number of return strokes, and which direction finders were used to locate the strike. This is done for the entire western United States. In 1985 a total of 1,913,602 strikes were recorded by the network. In 1986, 2,323,897 strikes were logged. For this study, the data was filtered into the grid block network illustrated in Figure 4. Each grid block is .5 degree latitude by .5 degree longitude and thus very close to the size used by Reap (1986) in the study of 1983-84 data. The grid is oriented differently than Reap's however. Our grid size was chosen for computer resource reasons and also to minimize the possibility of over-fitting the data based on current estimates of errors in specifying the location of a strike.

Since the BLM network was only operational for the months of April through October during 1985, those seven months of data were used for both 1985 and 1986 in constructing the preliminary climatologies. It appears that thunderstorms are quite rare in Idaho during the winter months of November to March. The authors do reserve the right to reverse their position on this after a few years of data have been assembled. In any event, current exclusion of these months may cause a slight underestimation of the "annual" frequencies found.

Figure 5 depicts the average annual number of lightning days based on the 1985 and 1986 BLM data. Stippling on the figure denotes generalized "mountainous" terrain. General valley areas are the Snake River Valley of southern Idaho and the Idaho Panhandle region. It should be noted that the Snake River Valley actually slopes from 600 meters near the western Idaho border to 1500 meters at its northeastern extension. Figure 5 indicates a general increase in the frequency of lightning days from west to east across the state. This trend appears to be definitely distorted by topography. Note that a relative minimum pushes eastward across the Snake River Valley while a relative maximum extends westward along the mountains near the state's southern border.

Comparison of Figure 5 with Figures 1 through 3 indicate two striking differences. First, this preliminary climatology suggests a significantly greater frequency than previously thought. Second, the maximum presented by Changery in Figure 3 appears to actually be further southeast and centered over the Yellowstone National Park area. The absolute maximum appears to occur over the mountains just east of the northern extension of the Snake River Valley. Interesting secondary maxima also appear to exist over the southwest Idaho mountains and the Wallowa Mountains in the northeast corner of Oregon.

As indicated by the values given for the total number of strikes recorded by the BLM network for the two years used, 1986 appears to have been a much more active thunderstorm year. This was true for our grid as well. In 1985 a total of 196,474 strikes fell within the grid. During 1986 the total was 262,809. Figure 6 depicts the number of lightning days for just 1985. Figure 7 is for 1986 alone. Comparison of these two figures indicates a difference in the number of days each year. However, the regions of relative maxima and minima coincide almost exactly. This lends strong credence to the hypothesis of terrain-induced convection. However, terrain height does not appear to be the dominant factor.

Figure 8 depicts the average number of hours with lightning strikes per year. Again the pattern coincides quite well with the pattern of lightning days across the state. When one looks at the average number of actual lightning strikes per year as shown in Figure 9, a significantly different pattern emerges. The region around Yellowstone which stood out as a center of maximum lightning days and hours now appears as a relative minimum. When considering total lightning strikes, the absolute maximum shifts to near the southeast corner of Idaho. Caution must be exercised in drawing conclusions from this, however, as operational experience has shown that occasionally a thunderstorm can become extremely active with regard to the amount of cloud-to-ground lightning it produces. The analysis does suggest that these highly active thunderstorms may be quite rare over some locations and show a preference for other areas.

### III. FREQUENCY OF LIGHTNING UNDER VARIOUS SYNOPTIC FLOW PATTERNS

Figure 10 portrays the two year average percent of days with lightning for the summer season of June, July and August. As can be seen, the summer season climatological probability of a day with lightning varies from less than 20 percent over much of western Idaho to 50 percent in the Upper Snake River Plain, the Yellowstone area and over the Teton Range.

To test whether the frequency of lightning days varies significantly under different synoptic scale flow patterns, the two year summer season data were stratified based on the 500 mb map types developed by Rasch and MacDonald (1975). The 1200 GMT 500 mb geopotential height analysis for each of the 184 sample days was assigned to the summer map type that it correlated with best. Figures 11 through 20 depict each of the 10 map types along with the corresponding frequency of occurrence of lightning during the period 1800 GMT to 0600 GMT (noon to midnight) on those days. Thus each figure gives the probability of lightning for each grid block under the accompanying synoptic scale mid-tropospheric flow pattern.

Figure 11 gives the Summer Type 1 flow pattern and corresponding probability of lightning. As with the data used to develop the map types, type 1 cases made up the largest percentage of the two year sample used in this study. Type 1 is a weak zonal pattern that

occurs over the western United States frequently during the summer months. Thus, as expected, the Type 1 frequencies are very similar to the seasonal mean frequencies shown in Figure 10. The probabilities over the southern portion of the state are slightly higher and those over the north a little lower than the mean.

Summer Type 2 probabilities presented in Figure 12 indicate a marked deviation from the means of Figure 10. Type 2 cases are characterized by the axis of the strongest upper tropospheric winds over or just south of southern Idaho. Thus dynamics associated with the jet stream may account for the much higher than normal probabilities over southern Idaho. The cold core areas of low pressure aloft associated with Type 2 cases often tracks across northern Idaho. This may explain the increased probabilities over the Panhandle.

Map Type 3, with an upper ridge axis over or just east of Idaho, was the second most prevalent pattern in both the data set used to develop the map types, and for 1985, 1986. The corresponding probability of lightning chart in Figure 13 is even closer to the seasonal climatological probability than Type 1. Main differences are more of a minimum in the Snake River Valley and a stronger maximum over the Wallowa Mountains of northeast Oregon.

Map Type 4 shown in Figure 14 is characterized by a strong upper ridge near 100 degrees west longitude and a fairly deep trough on or just off the West Coast. The jet stream axis is usually, but not always, north and west of Idaho. In some cases the upper flow over Idaho, especially the southern portion, is very weak. This pattern can be quite moist. The corresponding probability chart reflects this with a relatively high occurrence of lightning over the Snake River Valley.

Figure 15 shows Type 5 which is characterized by a split flow or a shearing trough over the state. The associated lightning probability chart suggests the troughs in the northern branch of the split are generally dry until they interact with the usually more moist weak southerly flow over southeast Idaho and western Wyoming.

Map Type 6, shown in Figure 16, is also characterized by a split flow. In this case the flow over Idaho is northwesterly. As expected, the corresponding lightning activity is quite low over most of Idaho under this pattern. In the light flow aloft over Montana, lightning episodes are more frequent.

The dominant feature for Type 7 days is a short wave ridge moving over Idaho in the wake of a trough. As shown in Figure 17, thunderstorm activity is well below the climatological frequency over the entire region for this type of flow pattern.

Probabilities are also quite low for Type 8 as shown on Figure 18. The glaring exception is the intriguing high frequency area over Yellowstone Park. The anomaly suggests that the terrain of that area may somehow react very favorably with short waves moving south in the northerly flow. There were only 17 Type 8 cases so it is not possible to say whether or not the anomaly is a stable one.

Map Type 9 in Figure 19 is an even rarer occurrence, comprising about one-half of one percent of all the cases used to construct the map types and about two percent of the 1985-1986 cases. The accompanying probability chart is only included for completeness. No real conclusions can be drawn from such a small sample size. However, as expected, the pattern does appear to produce considerable thunderstorm activity over Idaho.

Map Type 10 as depicted in Figure 20 is dominated by a broad ridge of high pressure over the western United States. The winds over Idaho are extremely light. Thus this map type may be an excellent starting point in investigating how mountainous terrain modifies the distribution of so-called "air-mass" thunderstorms. The associated lightning probability chart suggests that a few additional years of data may in fact provide some concrete insight into this problem. Note the numerous maxima and minima in Figure 20. These appear to generally correspond with significant terrain features. Equally important, if one compares Figure 20 with the more dynamic types 2 and 4, these "air-mass" centers generally tend to correspond to "dynamic" centers of the opposite sign.

#### IV. MANUALLY OBSERVED LIGHTNING ACTIVITY LEVEL

The National Fire Danger Rating System (NFDRS) implemented in 1972 is a computer model used by all federal and many state land management agencies to calculate current and forecast fire potential. In the western United States, lightning is the greatest single cause of forest fires. A critical element used to determine fire danger in the NFDRS is a lightning risk factor.

Prior to the introduction of the NFDRS, lightning forecasts were in probability terms. This approach made verification difficult. Studies of lightning and fire starts during Project Skyfire (1969) and later studies by Fuquay (1980) lead to the lightning activity level (LAL) index used by the NFDRS.

The LAL is a numerical rating which ranges from 1 to 6. It is keyed to the frequency and characteristics of the cloud-to-ground lightning observed or forecast for a specified area and time period. The scale is exponential in powers of 2. Beginning with LAL 2, each succeeding level through LAL 5 represents twice the cloud-to-ground lightning activity as the previous level. LAL 6 is a special and rare case in which high level dry thunderstorms create severe fire problems. LAL 1 indicates no lightning during the specified rating period.

Several schemes have been devised to help the fire weather observer and fire weather forecaster determine the appropriate LAL category. The Lightning Activity Guide (Deeming 1978) is the standard by which the LAL is presently calculated. The Guide has been divided into two sections. The first uses cloud descriptions and precipitation conditions to estimate the LAL category. The second uses lightning rates and amount estimates to arrive at a representative LAL category. The basic unit of area represented by a LAL is 2,500 square miles. This is accepted as the largest area that lightning activity can be effectively observed from a surface observation point such as a fire lookout. It roughly corresponds to a circle around the observation site with radius of about 45 km.

The observer subjectively assigns a LAL value. The ALDS network now provides the opportunity to compare these subjective reports with actual lightning occurrence. For this comparison 18 observation points in the Boise Fire Weather District were chosen. These 18 locations correspond to the established verification stations in each fire weather forecast zone as shown in Figure 21. Two summers (1985, 1986) of data were available for these comparisons.

Figure 22 shows the relative frequency distribution of LAL 2 through LAL 6 upon which the NFDRS is based. Also shown is the frequency distribution of the manually reported LAL for the 18 sites during the two year study period. As Figure 22 indicates, the manual LAL's were highly weighted toward the lower LAL categories. The frequency dropped off rapidly in the higher categories. This distribution is far from the frequency distribution the NFDRS expects, i.e., emphasis on the middle categories.

For each day, an ALDS LAL was assigned to each observation site. This was done by computing the total number of cloud-to-ground lightning strikes which fell within a 45 km radius of the observation site and then using the lightning density limits established for each LAL category by the NFDRS. For example, if 40 lightning strikes were recorded within 45 km of a station, a "true" LAL of 3 was assigned since the NFDRS range for LAL 3 is 11 to 50 strikes in a 24-hour period. Table 1 shows the percentage of correct manual observations as verified by the ALDS data. As shown, the frequency of correct categories during thunderstorm days is generally 50 percent or less. The only category which shows a high degree of skill is LAL 5. This is probably due to the fact that this category has the widest limits. Also, the station (Malad) which contained the bulk of LAL 5 cases was near the lightning maximum for the region as indicated in Figure 9. The LAL 1 category should be considered independently since it represents days with no thunderstorms. Table 1 indicates that observers totally missed 28 percent of the lightning occurrences the past two summers. The table also shows that only reported LAL categories 1 and 5 were generally representative of the actual lightning density that occurred.

A look at the means, standard deviations, and ranges of true lightning densities (ALDS data) corresponding to the manually reported LAL's is shown in Table 2. The means for each category are significantly higher than the established NFDRS limits (Table 3). Note also that the standard deviations and ranges are quite large and many overlap. Table 2 does indicate that a few observing sites were able to show skill in assigning LAL's which were representative of actual lightning density in a relative sense. Note, for example, the data for Stanley shows a general doubling of the lightning density mean value between LAL categories. The same is true for Mammoth and to some degree, Island Park.

Since the data from most observing sites showed very little correlation or apparent trends between the ALDS observed lightning densities and the manual reported densities (through the reported LAL), we investigated the possibility of adjusting the current NFDRS limits to improve the manually reported LAL verification.

The new limits shown in Table 3 we obtained by forcing the ALDS determined lightning densities for all 18 sites and all days into the NFDRS expected frequency distribution. As shown in Table 3 the data produced new limits which had lower bounds for all but categories 2 and 5. Also the ranges for categories 3 and 6 were greatly reduced while the limits for category 4 were significantly widened. When the manually reported LAL's were compared to the proposed new limits, the only category that experienced improvement was LAL category 4 as shown in Figure 23.

The frequency distribution of reported LAL's which the NFDRS expects to receive could be changed. This in turn might lead to better manually observed LAL verification scores by again utilizing the ALDS data to adjust the LAL category lightning density ranges. This "tampering" does not appear advisable, however, since it appears the problem lies not with the system, but with the manually reported LAL program.

## V. DISCUSSION

The actual percentage of cloud-to-ground lightning strikes detected by the BLM network is still a matter of controversy. The most accepted figures vary from 50 to 75 percent. In addition, not all thunderstorms produce cloud-to-ground strikes as they pass over an area. Thus, one would expect a lightning days climatology, as presented here, to be a conservative estimate of thunderstorm days. However, the preliminary climatology suggests that previous climatologies significantly underestimate the frequency of days with thunderstorms over much of Idaho. It must be kept in mind, however, that the preliminary climatology presented is based on only two years of data. In addition, the grid block size used yields an effective "observer radius" of about 27 km which is somewhat larger than the radius of most human observations which are the basis for conventional thunderstorm climatologies.

The use of map types as a means of stratifying summer days and developing departures from the climatological frequencies appears promising. This approach could prove very useful in forecasting since the output from numerical models which verifies the best is mid-tropospheric synoptic scale geopotential height patterns. There are insufficient cases for a number of the map types to consider the current associated probability maps statistically stable, however. It also appears that the inclusion of an atmospheric stability or moisture parameter is needed to elevate the map type approach to a viable short-term detailed forecasting tool. In addition, we are looking at hourly frequencies to determine if there are preferred areas of initial thunderstorm generation. We also intend to decrease the grid block size to one-quarter of a degree which will give an effective "observer radius" of about 14 km to see if this reveals smaller scale terrain influences.

The portion of the study dealing with lightning activity levels points to serious flaws in the manual observations used by the NFDRS. Clearly the observer's bias towards the lower LAL's and the tendency for the observed categories to have a higher density singles out the current misunderstanding and shortcomings of the observational system. In the near future, the Boise Fire Weather Office will be developing an automatic LAL verification program using the lightning detection data and the current NFDRS limits. Since the LAL guide was developed before the advent of this more comprehensive observational tool, further climatological studies will be needed before a readjustment can be made to the NFDRS frequency distributions and lightning density limits.

## VI. REFERENCES

- Changery, M.J., 1981: National Thunderstorm Frequencies for the Contiguous United States. U.S. Nuclear Regulatory Commission Report, NUREG/CR-2252.
- Climate and Man, 1941: Yearbook of Agriculture, U.S. Government Printing Office, p. 729.
- Critchfield, H.J., 1966: General Climatology, Prentice-Hall, Inc., Englewood Cliffs, New Jersey, p. 185.
- Deeming, J.E., R.E. Burgan, and J.D. Cohen, 1977: The National Fire Danger Rating System - 1978. USDA Forest Service Gen. Tech. Rep. INT-39, pp. 36-40.
- Fuquay, D.M., and Robert G. Baughman, 1969: Project Skyfire Lightning Research. Final rep. to Natl. Sci. Found., Grant GP-2617.
- Fuquay, D.M. 1980: Forecasting Lightning Activity level and Associated Weather. USDA Forest Service Res. Pup. INT-244.

- Rasch, G.E. and A.E. MacDonald, 1975: Map Tyle Precipitation Probabilities for the Western Region. NOAA Technical Memorandum NWS-WR-96.
- Rasch, G.E. and M. Mathewson, 1984: Collection and use of Lightning Strike Data in the Western U.S. During Summer 1983. NOAA Technical Memorandum NWR-WR-184.
- Reap, R.M. 1986: Evaluation of Cloud-to-Ground Lightning Data from the Western United States for the 1983-1984 Summer Seasons, J. Climate Appl. Meteor., 25, pp. 785-799.




Figure 1. Average Annual Number of Days with Thunderstorms (from "Climate and Man", 1941)




Figure 2. Average Annual Number of Days with Thunderstorms (from Critchfield, 1966).





Figure 3. Average Annual Number of Days with Thunderstorms (from Changery, 1981).

Figure 4. Grid block (one-half by one-half degree) array for which lightning data is archived and analyzed.






Figure 5. Average Annual Number of Days with Lightning Strikes based on two years (1985, 1986) of data.




Figure 6. Number of Days with Lightning Strikes during 1985.




Figure 7. Number of Days with Lightning Strikes during 1986.




Figure 8. Average Annual Number of Hours with Lightning Strikes based on two years (1985, 1986) of data.




Figure 9. Average Annual Number of Lightning Strikes. Isoceraunics are labeled as 100's of strikes. Maximum value in southeast Idaho is 2644. Based on two years (1985,1986) of data.




Figure 10. Average Annual Percent of Summer (June, July, August) Days with Lightning Strikes.

SUMMER TYPE I

00Z July 19, 1963

663 Cases





Figure 11. Summer Map Type 1 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 41 cases.

SUMMER TYPE 2

00Z June 22, 1963  
78 Cases

22





Figure 12. Summer Map Type 2 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 12 cases.

SUMMER TYPE 3

12Z June 3, 1964

135 Cases





Figure 13. Summer Map Type 3 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 28 cases.

SUMMER TYPE 4

00Z August 26, 1968  
86 Cases





Figure 14. Summer Map Type 4 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 28 cases.

SUMMER TYPE 5  
12Z June 8, 1963  
42 Cases





Figure 15. Summer Map Type 5 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 13 cases.

SUMMER TYPE 6

12Z August 9, 1966  
60 Cases





Figure 16. Summer Map Type 6 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 10 cases.

SUMMER TYPE 7

12Z June 10, 1968

52 Cases





Figure 17. Summer Map Type 7 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 12 cases.

SUMMER TYPE 8

00Z June 23, 1967  
76 Cases





Figure 18. Summer Map Type 8 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 17 cases.

SUMMER TYPE 9

00Z June 2, 1967  
7 Cases





Figure 19. Summer Map Type 9 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 4 cases.

SUMMER TYPE 10

12Z July 12, 1964

9 Cases





Figure 20. Summer Map Type 10 and Corresponding Percent Probabilities of Lightning during the period noon to midnight mdt. Probabilities based on 20 cases.




Figure 21. Map of verification stations.




Figure 22. NFDRS Relative Frequency Distribution versus Observed Frequency Distribution.




Figure 23. Frequency Distribution of Correct Observations Using NFDRS Limits versus Correct Observations Using Adjusted Limits.

TABLE 1

Frequency of Correct Categories Verified Against NFDRS  
Cloud to Ground Lightning Density Limits (Fuquay)  
1985-1986 Thunderstorm Season

STATION	LAL 1	LAL 2	LAL 3	LAL 4	LAL 5	LAL 6
McCall	84% (192)	18% (11)	10% (10)	50% (2)	100% (2)	NR (0)
Krassel	81% (171)	22% (9)	100% (4)	NR (0)	67% (3)	NR (0)
Garden Valley	84% (187)	32% (19)	33% (10)	0% (3)	NR (0)	0% (1)
Willow Creek	80% (190)	36% (14)	75% (4)	0% (2)	50% (2)	NR (0)
Indianola	73% (141)	26% (23)	22% (9)	NR (0)	100% (1)	NR (0)
Challis	66% (181)	25% (8)	25% (8)	40% (5)	100% (2)	100% (1)
Stanley	79% (161)	43% (14)	44% (9)	14% (7)	100% (3)	NR (0)
Burns Junction	89% (65)	26% (27)	0% (10)	NR (0)	100% (1)	0% (1)
Boise	93% (193)	33% (21)	0% (5)	0% (6)	0% (1)	67% (3)
Notch Butte	64% (143)	50% (2)	25% (4)	100% (1)	NR (0)	NR (0)
Dubois	62% (169)	31% (32)	10% (10)	25% (12)	0% (1)	NR (0)
Island Park	48% (191)	28% (18)	0% (6)	0% (5)	75% (4)	NR (0)
Rock Creek	69% (132)	0% (15)	0% (4)	34% (3)	0% (1)	NR (0)
Malad	76% (84)	12% (17)	40% (10)	22% (9)	76% (25)	50% (4)
Big Piney	56% (134)	26% (19)	22% (18)	0% (2)	83% (6)	NR (0)
Black Rock	66% (146)	14% (28)	36% (11)	0% (4)	100% (1)	50% (2)
Mammoth	61% (143)	25% (36)	31% (26)	11% (9)	25% (4)	100% (1)
Rock Springs	55% (86)	30% (63)	37% (30)	50% (2)	100% (1)	33% (3)
<b>TOTALS</b>	72% (2,709)	26% (376)	29% (188)	18% (72)	72% (58)	50% (16)

Total number of events in parenthesis

TABLE 2

Lightning Density Mean,  
Standard Deviation and Range  
for Man Observed Thunderstorm Days  
by LAL Category

Station		LAL 2	LAL 3	LAL 4	LAL 5	LAL 6
McCall	Mean	91	42	44	276	NR
	S.D.	101.0	43.2	—	—	
	Range	0-277	0-127	6-82	206-346	
Krassell	Mean	52	28	NA	102	NR
	S.D.	66.5	8.7		76.9	
	Range	0-214	23-41		15-160	
Garden Valley	Mean	11	46	245	NR	199
	S.D.	13.8	73.8	216.8		—
	Range	0-47	0-242	46-476		—
Willow Creek	Mean	45	18	149	79	NR
	S.D.	89.8	12.4	—	—	
	Range	0-334	4-34	38-260	24-134	
Indianola	Mean	78	147	NR	322	NR
	S.D.	83.4	154.0		—	
	Range	0-302	3-431		—	
Challis	Mean	47	149	123	181	24
	S.D.	47.5	153.2	153.3	—	—
	Range	1-151	2-389	19-392	134-227	—
Stanley	Mean	21	42	85	137	NR
	S.D.	22.3	36.4	101.3	8.4	
	Range	0-59	1-99	4-267	132-147	
Burns Junction	Mean	22	79	NR	118	0
	S.D.	72.6	247.3		—	—
	Range	0-360	0-783		—	—
Boise	Mean	12	4	86	25	39
	S.D.	12.8	3.2	75.2	—	23.5
	Range	0-50	0-8	10-197	—	16-63
North Butte	Mean	72	126	85	NR	NR
	S.D.	100.4	83.8	—		
	Range	1-143	19-198	—		
Dubois	Mean	75	216	191	25	NR
	S.D.	121.7	179.6	186.1	—	
	Range	0-505	48-574	1-324	—	

STATION		LAL 2	LAL 3	LAL 4	LAL 5	LAL 6
Island Park	Mean	51	174	327	224	NR
	S.D.	49.9	125.7	268.1	220.7	
	Range	1-149	84-395	114-753	51-545	
Rock Creek	Mean	129	233	133	78	NR
	S.D.	222.5	156.8	75.8	--	
	Range	0-876	136-465	68-216	--	
Malad	Mean	59	46	208	426	46
	S.D.	72.6	67.3	333.1	417.1	37.7
	Range	0-206	0-190	1-1,049	14-1,659	12-82
Big Piney	Mean	130	193	335	282	NR
	S.D.	169.5	216.8	--	158	
	Range	0-476	0-709	12-657	72-478	
Black Rock	Mean	115	58	224	257	103
	S.D.	171.2	42.7	212.1	--	--
	Range	0-815	0-109	4-507	--	17-188
Mammoth	Mean	49	98	205	281	23
	S.D.	77.3	96.9	138.2	384.2	--
	Range	0-366	0-392	17-466	73-857	--
Rock Springs	Mean	84	94	106	272	52
	S.D.	113.2	90.7	--	--	71.4
	Range	0-391	4-337	60-152	--	8-134
<b>TOTALS</b>	Mean	72	104	177	298	57
	S.D.	114.1	136.1	197.2	321.9	64.1
	Range	0-876	0-783	1-1,049	14-1,659	8-199

TABLE 3

Number of Cloud to Ground  
Strikes within 2,500 mi<sup>2</sup> Area  
Per LAL Category

LAL Category	1	2	3	4	5	6
NFDRS (Fuquay) Limits	No lightning	1-10	11-50	51-100	>100	11-50
New Limits To Obtain NFDRS Frequency Distribution	No lightning	1	2-15	16-118	>118	2-15
NFDRS Relative Frequency of Thunderstorm Days per Category	No lightning	10%	35%	35%	18%	2%

- 139 Aids for Forecasting Minimum Temperature in the Wenatchee Frost District. Robert S. Robinson, April 1979. (PB298339/AS)
- 140 Influence of Cloudiness on Summertime Temperatures in the Eastern Washington Fire Weather district. James Holcomb, April 1979. (PB298674/AS)
- 141 Comparison of LFM and MFM Precipitation Guidance for Nevada During Doreen. Christopher Hill, April 1979. (PB298613/AS)
- 142 The Usefulness of Data from Mountaintop Fire Lookout Stations in Determining Atmospheric Stability. Jonathan W. Corey, April 1979. (PB298899/AS)
- 143 The Depth of the Marine Layer at San Diego as Related to Subsequent Cool Season Precipitation Episodes in Arizona. Ira S. Brenner, May 1979. (PB298817/AS)
- 144 Arizona Cool Season Climatological Surface Wind and Pressure Gradient Study. Ira S. Brenner, May 1979. (PB298900/AS)
- 146 The BART Experiment. Morris S. Webb, October 1979. (PB80 155112)
- 147 Occurrence and Distribution of Flash Floods in the Western Region. Thomas L. Dietrich, December 1979. (PB80 160344)
- 149 Misinterpretations of Precipitation Probability Forecasts. Allan H. Murphy, Sarah Lichtenstein, Baruch Fischhoff, and Robert L. Winkler, February 1980. (PB80 174576)
- 150 Annual Data and Verification Tabulation - Eastern and Central North Pacific Tropical Storms and Hurricanes 1979. Emil B. Gunther and Staff, EPHC, April 1980. (PB80 220486)
- 151 NMC Model Performance in the Northeast Pacific. James E. Overland, PMEL-ERL, April 1980. (PB80 196033)
- 152 Climate of Salt Lake City, Utah. Wilbur E. Figgins, Third Revision January 1987. (PB87 157194/AS)
- 153 An Automatic Lightning Detection System in Northern California. James E. Rea and Chris E. Fontana, June 1980. (PB80 225592)
- 154 Regression Equation for the Peak Wind Gust 6 to 12 Hours in Advance at Great Falls During Strong Downslope Wind Storms. Michael J. Oard, July 1980. (PB91 108367)
- 155 A Raininess Index for the Arizona Monsoon. John H. Ten Harkel, July 1980. (PB81 106494)
- 156 The Effects of Terrain Distribution on Summer Thunderstorm Activity at Reno, Nevada. Christopher Dean Hill, July 1980. (PB81 102501)
- 157 An Operational Evaluation of the Scofield/Oliver Technique for Estimating Precipitation Rates from Satellite Imagery. Richard Ochoa, August 1980. (PB81 108227)
- 158 Hydrology Practicum. Thomas Dietrich, September 1980. (PB81 134033)
- 159 Tropical Cyclone Effects on California. Arnold Court, October 1980. (PB81 133779)
- 160 Eastern North Pacific Tropical Cyclone Occurrences During Intraseasonal Periods. Preston W. Leftwich and Gail M. Brown, February 1981. (PB81 205494)
- 161 Solar Radiation as a Sole Source of Energy for Photovoltaics in Las Vegas, Nevada, for July and December. Darryl Randerson, April 1981. (PB81 224503)
- 162 A Systems Approach to Real-Time Runoff Analysis with a Deterministic Rainfall-Runoff Model. Robert J.C. Burnash and R. Larry Ferral, April 1981. (PB81 224495)
- 163 A Comparison of Two Methods for Forecasting Thunderstorms at Luke Air Force Base, Arizona. LTC Keith R. Cooley, April 1981. (PB81 225393)
- 164 An Objective Aid for Forecasting Afternoon Relative Humidity Along the Washington Cascade East Slopes. Robert S. Robinson, April 1981. (PB81 23078)
- 165 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1980. Emil B. Gunther and Staff, May 1981. (PB82 230336)
- 166 Preliminary Estimates of Wind Power Potential at the Nevada Test Site. Howard G. Booth, June 1981. (PB82 127036)
- 167 ARAP User's Guide. Mark Mathewson, July 1981, revised September 1981. (PB82 195783)
- 168 Forecasting the Onset of Coastal Gales Off Washington-Oregon. John R. Zimmerman and William D. Burton, August 1981. (PB82 127051)
- 169 A Statistical-Dynamical Model for Prediction of Tropical Cyclone Motion in the Eastern North Pacific Ocean. Preston W. Leftwich, Jr., October 1981. (PB82 195298)
- 170 An Enhanced Plotter for Surface Airways Observations. Andrew J. Spry and Jeffrey L. Anderson, October 1981. (PB82 153883)
- 171 Verification of 72-Hour 500-MB Map-Type Predictions. R.F. Quiring, November 1981. (PB82 158098)
- 172 Forecasting Heavy Snow at Wenatchee, Washington. James W. Holcomb, December 1981. (PB82 177783)
- 173 Central San Joaquin Valley Type Maps. Thomas R. Crossan, December 1981. (PB82 196064)
- 174 ARAP Test Results. Mark A. Mathewson, December 1981. (PB82 198103)
- 176 Approximations to the Peak Surface Wind Gusts from Desert Thunderstorms. Darryl Randerson, June 1982. (PB82 253089)
- 177 Climate of Phoenix, Arizona. Robert J. Schmidl, April 1969 (revised December 1986). (PB87 142063/AS)
- 178 Annual Data and Verification Tabulation, Eastern North Pacific Tropical Storms and Hurricanes 1982. E.B. Gunther, June 1983. (PB85 106078)
- 179 Stratified Maximum Temperature Relationships Between Sixteen Zone Stations in Arizona and Respective Key Stations. Ira S. Brenner, June 1983. (PB83 249904)
- 180 Standard Hydrologic Exchange Format (SHEF) Version I. Phillip A. Pasteries, Vernon C. Bissel, David G. Bennett, August 1983. (PB85 106052)
- 181 Quantitative and Spatial Distribution of Winter Precipitation along Utah's Wasatch Front. Lawrence B. Dunn, August 1983. (PB85 106912)
- 182 500 Millibar Sign Frequency Teleconnection Charts - Winter. Lawrence B. Dunn, December 1983. (PB85 106276)
- 183 500 Millibar Sign Frequency Teleconnection Charts - Spring. Lawrence B. Dunn, January 1984. (PB85 111367)
- 184 Collection and Use of Lightning Strike Data in the Western U.S. During Summer 1983. Glenn Rasch and Mark Mathewson, February 1984. (PB85 110534)
- 185 500 Millibar Sign Frequency Teleconnection Charts - Summer. Lawrence B. Dunn, March 1984. (PB85 111359)
- 186 Annual Data and Verification Tabulation eastern North Pacific Tropical Storms and Hurricanes 1983. E.B. Gunther, March 1984. (PB85 109635)
- 187 500 Millibar Sign Frequency Teleconnection Charts - Fall. Lawrence B. Dunn, May 1984. (PB85 110930)
- 188 The Use and Interpretation of Isentropic Analyses. Jeffrey L. Anderson, October 1984. (PB85 132694)
- 189 Annual Data & Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1984. E.B. Gunther and R.L. Cross, April 1985. (PB85 187887AS)
- 190 Great Salt Lake Effect Snowfall: Some Notes and An Example. David M. Carpenter, October 1985. (PB86 119153/AS)
- 191 Large Scale Patterns Associated with Major Freeze Episodes in the Agricultural Southwest. Ronald S. Hamilton and Glenn R. Luskky, December 1985. (PB86 144474AS)
- 192 NWR Voice Synthesis Project: Phase I. Glen W. Sampson, January 1986. (PB86 145604/AS)
- 193 The MCC - An Overview and Case Study on Its Impact in the Western United States. Glenn R. Luskky, March 1986. (PB86 170651/AS)
- 194 Annual Data and Verification Tabulation Eastern North Pacific Tropical Storms and Hurricanes 1985. E.B. Gunther and R.L. Cross, March 1986. (PB86 170941/AS)
- 195 Rapid Interpretation Guidelines. Roger G. Pappas, March 1986. (PB86 177680/AS)
- 196 A Mesoscale Convective Complex Type Storm over the Desert Southwest. Darryl Randerson, April 1986. (PB86 190998/AS)
- 197 The Effects of Eastern North Pacific Tropical Cyclones on the Southwestern United States. Walter Smith, August 1986. (PB87 106258AS)

## NOAA SCIENTIFIC AND TECHNICAL PUBLICATIONS

*The National Oceanic and Atmospheric Administration* was established as part of the Department of Commerce on October 3, 1970. The mission responsibilities of NOAA are to assess the socioeconomic impact of natural and technological changes in the environment and to monitor and predict the state of the solid Earth, the oceans and their living resources, the atmosphere, and the space environment of the Earth.

The major components of NOAA regularly produce various types of scientific and technical information in the following kinds of publications:

**PROFESSIONAL PAPERS** — Important definitive research results, major techniques, and special investigations.

**CONTRACT AND GRANT REPORTS** — Reports prepared by contractors or grantees under NOAA sponsorship.

**ATLAS** — Presentation of analyzed data generally in the form of maps showing distribution of rainfall, chemical and physical conditions of oceans and atmosphere, distribution of fishes and marine mammals, ionospheric conditions, etc.

**TECHNICAL SERVICE PUBLICATIONS** — Reports containing data, observations, instructions, etc. A partial listing includes data serials; prediction and outlook periodicals; technical manuals, training papers, planning reports, and information serials; and miscellaneous technical publications.

**TECHNICAL REPORTS** — Journal quality with extensive details, mathematical developments, or data listings.

**TECHNICAL MEMORANDUMS** — Reports of preliminary, partial, or negative research or technology results, interim instructions, and the like.



*Information on availability of NOAA publications can be obtained from:*

**ENVIRONMENTAL SCIENCE INFORMATION CENTER (D822)  
ENVIRONMENTAL DATA AND INFORMATION SERVICE  
NATIONAL OCEANIC AND ATMOSPHERIC ADMINISTRATION  
U.S. DEPARTMENT OF COMMERCE**

**6009 Executive Boulevard  
Rockville, MD 20852**